

Photometric variability of 14 PMS stars in the NGC 7000/IC 5070 complex

Sunay I. Ibryamov^{1,2} and Evgeni H. Semkov¹

¹ Institute of Astronomy and NAO, Bulgarian Academy of Sciences, BG-1784 Sofia

² Department of Theoretical and Applied Physics, University of Shumen, BG-9712 Shumen
sibryamov@astro.bas.bg

(Submitted on; Accepted on)

Abstract. New photometric data from CCD multicolour *BVRI* observations of 14 pre-main sequence stars during the period from 2013 April to 2015 September are presented. The studied objects are located in the field of 'Gulf of Mexico' in the NGC 7000/IC 5070 star-forming complex. The stars from our study exhibit different types of photometric variability in all optical passbands. Using our long-term observations and data published by other authors, we tried to define the reasons for the observed brightness variations. On the basis of our new data previously unknown periodicity in the light curve of the star LkH α 189 (2.45 days) was registered.

Key words: stars: pre-main sequence, stars: variables: T Tauri, star: individual: V521 Cyg, V752 Cyg, V1538 Cyg, V1539 Cyg, V1716 Cyg, V1957 Cyg, V2051 Cyg, LkH α 186, LkH α 187, LkH α 189, LkH α 191, [KW97] 53-11, [KW97] 53-23, [KW97] 53-36

1. Introduction

Studies of pre-main sequence (PMS) stars give an opportunity to understand the early stages of stellar evolution and to test stellar evolution scenarios. Depending on their initial mass, PMS stars pass through different periods of stellar activity. The most prominent manifestations of this activity are changes in the star's brightness with various periods and amplitudes.

PMS stars are rare among field stars, because during the PMS evolutionary phase the stars spend less than 1% of its life. PMS stars are separated into two main types, low mass ($M \leq 2M_{\odot}$) T Tauri stars (TTS) and the more massive ($2M_{\odot} \leq M \leq 8M_{\odot}$) Herbig Ae/Be stars (HAEBES). Both TTS and HAEBES still contracting towards the main sequence.

TTS are associated with dark nebulae, molecular clouds, obscured regions, and are grouped in T associations. The main characteristics of TTS are their strong irregular photometric variability and emission line spectra (Joy 1945). TTS are divided into two sub-classes: classical T Tauri stars (CTTS) and weak-line T Tauri stars (WTTS). Most of the features that characterize each sub-class suggest that accretion disks surround CTTS whereas they must have almost disappeared (at least their inner parts) in WTTS (see Bertout 1989; Ménard & Bertout 1999).

According to Herbst et al. (2007), variability of CTTS can be due to rotating hot spots on the stellar surface. CTTS also often shows irregular variations with large photometric amplitudes up to 2-3 mag, associated with high variable accretion from the circumstellar disk onto the stellar surface. The variability of WTTS is due to cool spots or groups of spots on the stellar surface. The photometric amplitudes of this variability are about 0.03-0.3 mag, but in extreme cases reach to 0.8 mag in the *V*-band. The variability of WTTS can be due to flare-like variations in *B*- and *U*-band also. Flares are random with different sized amplitudes, as there is no periodicity. Usually, the rise of brightness is quick and occurs over short time scales.

The large amplitude drops in the brightness in the light curves are commonly observed in the early types (G, F) of TTS and HAEBS. These stars show non-periodic deep Algol-type minima and photometric amplitudes up to 2.8 mag in *V*-band. The observed drops in the brightness last from days to some weeks and are caused by variable extinction from circumstellar dust or clouds (Voshchinnikov 1989; Grinin et al. 1991; Herbst et al. 2007). This group of PMS stars with intermediate mass is known as UXors and their prototype is the star UX Orionis. In very deep minima, the colour indices of UXors often becomes bluer (so called 'blueing effect' or 'colour reverse') (see Bibó & Thé 1990). It is generally accepted that the origin of observed drops in the brightness and the blueing effect are due to variations of the column density of dust in the line of sight to the star (Dullemond et al. 2003).

The stars from our study are located in the dense molecular cloud L935, known as 'Gulf of Mexico' in the NGC 7000/IC 5070 complex. The NGC 7000/IC 5070 complex are thought to be parts of a single large HII region W80. The distance to this region, determined by Laugalys & Straizys (2002) is 600 pc. 'Gulf of Mexico' is a region with active star formation and contains many PMS objects (see Armond et al. 2011; Findeisen et al. 2013; Bally et al. 2014; Poljančič Beljan et al. 2014). The PMS stars included in the present study are revealed in previous studies of the 'Gulf of Mexico' as H α emission stars, flare stars from UV Cet type, TTS or HAEBS.

The present paper is a part of our long-term multicolour photometric study of the PMS stars in the region of 'Gulf of Mexico'. The results from our previous studies of PMS stars in this field have been published in Semkov et al. (2010, 2012, 2014), Poljančič Beljan et al. (2014), Ibryamov et al. (2015a), and Ibryamov et al. (2015b). Section 2 gives information about telescopes and cameras used and data reduction. Section 3 describes the obtained results and their interpretation.

2. Observations and Data reduction

The presented photometric data were carried out in the period from 2013 April to 2015 September. The observations were performed with four telescopes: 2-m Ritchey-Chrétien-Coudé (RCC), the 50/70-cm Schmidt and the 60-cm Cassegrain telescopes of the Rozhen National Astronomical Observatory in Bulgaria and the 1.3-m Ritchey-Chrétien (RC) telescope of the Skinakas Observatory³ of the University of Crete in Greece. The total number of the nights used for observations is 84. A log of the observing periods and telescopes used is given in Table 1.

The observations were performed with four different types of CCD cameras: VersArray 1300B at the 2-m RCC telescope, ANDOR DZ436-BV at the 1.3-m RC telescope, FLI PL16803 at the 50/70-cm Schmidt telescope, and FLI PL09000 at the 60-cm Cassegrain telescope. The technical parameters and specifications for the cameras used, observational procedure, and data reduction process are given in Ibryamov et al. (2015b). The photometric limit of our data obtained with the 2-m RCC and the 1.3-m RC telescopes in *V*-

³ Skinakas Observatory is a collaborative project of the University of Crete, the Foundation for Research and Technology, Greece, and the Max-Planck-Institut für Extraterrestrische Physik, Germany.

Table 1. Photometric CCD observations of the field of 'Gulf of Mexico' with the different telescopes

Telescope	Nr of observations	Time span
2-m RCC	23	04.07.2013 – 06.09.2015
1.3-m RC	4	17.09.2013 – 12.08.2015
50/70-cm Schmidt	41	10.04.2013 – 03.09.2015
60-cm Cassegrain	16	15.05.2013 – 21.07.2014

and B -band is about 20.5 mag and for the data collected with the 50/70-cm Schmidt and the 60-cm Cassegrain telescopes is about 19.5 mag.

All frames were taken through a standard Johnson-Cousins (BVR_cI_c) set of filters. All data were analyzed using the same aperture, which was chosen to have a $4''$ radius, while the background annulus was taken from $9''$ to $14''$. As a reference, the $BVRI$ comparison sequence reported in Semkov et al. (2010) was used. The mean value of the errors in the reported magnitudes are 0.01-0.02 mag for I - and R -band data, 0.02-0.06 mag for V -band data, and 0.02-0.09 mag for B -band data.

3. Results and Discussion

The photometric results for 14 stars obtained in our study and their interpretation are presented in the section. All stars exhibit photometric variability in $BVRI$ optical passbands. Our data are a continuation of the long-term photometric investigations of these stars published in Poljančič Beljan et al. (2014) and Ibryamov et al. (2015a).

The data for the studied objects are presented in Table 2. Star identifiers used in this paper are marked in boldface. Figure 1 shows a three-colour images of the field of 'Gulf of Mexico', where the positions of the stars from our study and the recently erupting FUor star V2493 Cyg (see Semkov et al. 2010, 2012, 2014; Miller et al. 2011) are marked. The images were obtained with the 50/70-cm Schmidt telescope of Rozhen NAO on 2012 September 04 (the left image) and on 2014 August 19 (the right image).

The stars from our study shows bursts and/or fades events in their light curves. These events are defined as follows. In the case that the star most of the time spends at high light during our study, the declines in the brightness are interpreted as fades. Respectively, if the star most of the time spends at a low light, the observed increases in the brightness are interpreted as bursts.




Fig. 1. Three-colour images of the field of 'Gulf of Mexico'. The positions of the stars from our study and the recently erupting FUor star V2493 Cyg are marked. The images were obtained with the 50/70-cm Schmidt telescope of Rozhen NAO on 2012 September 04 (the left image) and on 2014 August 19 (the right image).

The JHK_s 2MASS magnitudes of the studied stars were used to construct the two-colour diagram and to identify stars with infrared excess, indicating the presence of a disk. Fig. 2 shows the location of main sequence (brown belt) and giant stars (orange belt) from Bessell & Brett (1988), and the classical T Tauri stars location (green belt) from Meyer et al. (1997). A correction to the 2MASS photometric system was performed following the prescription of Carpenter (2001). The three parallel dotted lines show the direction of the interstellar reddening vectors determined for the 'Gulf of Mexico' region by Straizys et al. (2008). In Fig. 2 the stars are designated using the numbers from Table 2. The stars lying to the right of the middle vector indicate the presence of a infrared excess. Due to the photometric variability, the positions of stars in Fig. 2 may vary around the displayed values.

Table 3 contain the mean photometrical magnitudes and colour indices for the stars from our study.

Table 2. Designations and coordinates of the stars from our study

Nr	GCVS ¹	HBC ²	[KW97] ³	LkH α ⁴	2MASS ID ⁵	$RA_{J2000.0}$	$Dec_{J2000.0}$
1	V521 Cyg	299	53-26	188	J20582381+4353114	20 58 23.81	+43 53 11.45
2	V752 Cyg		53-30		J20583359+4403354	20 58 33.60	+44 03 35.50
3	V1538 Cyg				J20575750+4350089	20 57 57.51	+43 50 08.95
4	V1539 Cyg	720	53-9	185	J20575986+4353260	20 57 59.86	+43 53 26.00
5	V1716 Cyg				J20580611+4353011	20 58 06.12	+43 53 01.17
6	V1957 Cyg				J20575651+4352362	20 57 56.52	+43 52 36.29
7	V2051 Cyg				J20574880+4350236	20 57 48.80	+43 50 23.60
8		723	53-24	186	J20581961+4353545	20 58 19.62	+43 53 54.52
9		724	53-25	187	J20582154+4353449	20 58 21.54	+43 53 44.96
10		725	53-27	189	J20582400+4353546	20 58 24.01	+43 53 54.61
11		301	53-42	191	J20590581+4357030	20 59 05.82	+43 57 03.08
12			53-11			20 58 03.00	+43 50 42.00
13			53-23		J20581951+4353449	20 58 19.52	+43 53 45.00
14			53-36		J20584756+4402574	20 58 47.52	+44 02 59.29

¹General Catalogue of Variable Stars (Samus et al. 2009); ²Herbig & Bell (1988); ³Kohoutek & Wehmeyer (1997); ⁴Herbig (1958); ⁵2Micron All-Sky Survey (Skrutskie et al. 2006).

3.1. V521 Cyg

Variability of V521 Cyg was discovered and the star was classified as a RW Aurigae type by Hoffmeister (1949). The author reported that the photographic magnitude of the star vary from 14.30 to 16.60 mag. The star is included in the list of H α emission-line star published in Herbig (1958). The author determined the photographic magnitude of the star $m_{pg}=15.00$ mag and its spectral type K0.

Filin (1974) presented observations of V521 Cyg obtained during the period 1969–1971 in system close to B . The author registered slowly fluctuations in the star's brightness in the interval 14.50–15.10 mag, as these fluctuations are characterized with deep drops in the brightness of V521 Cyg from the




Fig. 2. The $J - H$ versus $H - K_s$ diagram for the stars detected in all 3 bands in 2MASS catalogue.

Table 3. Mean photometrical magnitudes and colour indices for the stars from our study

Star	\bar{I}	\bar{R}	\bar{V}	\bar{B}	$\overline{V-I}$	$\overline{V-R}$	$\overline{B-V}$
V521 Cyg	12.11	12.93	13.76	15.03	1.66	0.84	1.26
V752 Cyg	14.98	15.89	16.61	17.48	1.63	0.72	0.87
V1538 Cyg	15.00	16.04	17.00	18.38	2.00	0.95	1.38
V1539 Cyg	13.51	14.51	15.55	16.90	2.04	1.04	1.35
V1716 Cyg	15.01	16.36	17.51	19.07	2.51	1.16	1.56
V1957 Cyg	14.57	15.43	16.26	17.50	1.70	0.83	1.24
V2051 Cyg	13.93	15.56	16.61	18.12	2.68	1.05	1.52
LkH α 186	14.60	16.07	17.38	19.00	2.77	1.31	1.63
LkH α 187	14.72	16.23	17.61	19.26	2.90	1.38	1.67
LkH α 189	13.97	15.29	16.54	18.21	2.57	1.25	1.67
LkH α 191	11.58	12.19	12.87	13.98	1.29	0.68	1.11
[KW97] 53-11*	16.18	18.65	-	-	-	-	-
[KW97] 53-23	14.99	16.92	18.55	20.33	3.57	1.66	2.01
[KW97] 53-36	11.63	11.88	12.28	13.05	0.65	0.40	0.78

* $\overline{R-I}=2.54$.

Algol type. Chavarría-K. et al. (1989) observed the star in $uvby-\beta$ photometric system and concluded that it is PMS star.

Herbst & Shevchenko (1999) observed V521 Cyg during the period 1986–1997 and determined the amplitude of its photometric activity $\Delta V=1.77$ mag, average stellar magnitude $\bar{V}=13.74$ mag, and colour indices $U-B=0.58$, $B-V=1.25$ and $V-R=1.18$ mag. Mitskevich & Pavlenko (2001) presented the light curve in V -band of the star for the period 1999 July–October. It is seen that for about 20 days (in 1999 August), the star faded by 0.85 mag. The star spend about 5 days in minimum light state whereafter in next 70 days the star’s brightness gradually returned to the previous maximum level. The authors suggest that the reason for the decline in brightness of V521 Cyg is probably obscuration from circumstellar clouds of dust. In the presented colour indices $V-I$ and $V-R$ of the star is not observed bluing effect, because the registered minimum from Mitskevich & Pavlenko (2001) is relatively shallow.

Grankin et al. (2007) conducted long-term study of V521 Cyg during the period 1986 June–2003 September. The authors presented the light curve in V -band of the star with brightness variations from 13.37 to 14.70 mag. Non-linear dependence of $V-R$ colour index versus stellar V magnitude is present in Grankin et al. (2007).

In $J-H/H-K_s$ diagram V521 Cyg is situated in the upper right-hand corner, close to the intrinsic T Tauri line in the CTTS location (Laugalys et al. 2006). Poljančič Beljan et al. (2014) determined the period of V521 Cyg, which is found to be 503 days. The cause of the period found is likely precession of the circumstellar disk or eclipses from a second component or by clouds of gas and dust orbiting the star.

The $BVRI$ light curves of V521 Cyg from all our CCD observations (Poljančič Beljan et al. 2014; Ibryamov 2015a; and the present paper) are shown in Figure 3. In the figure, circles denote CCD photometric data obtained with the 2-m RCC telescope, diamonds – the photometric data collected with the 1.3-m telescope, triangles – the photometric data acquired with the 50/70-cm Schmidt telescope, and squares – the photometric data taken with the 60-cm Cassegrain telescope. The results of our recent multicolour CCD observations of V521 Cyg are summarized in Table 4⁴. The columns contains data (DD/MM/YYYY format) and Julian data (J.D.) of the observations, $IRVB$ magnitudes of the star, telescope and CCD camera used.

The brightness of V521 Cyg during all period of our observations 1997–2015 vary in the range 11.78–12.85 mag for I -band, 12.55–14.05 mag for R -band, 13.32–14.89 mag for V -band, and 14.49–16.69 mag in B -band. The observed amplitudes are $\Delta I=1.07$ mag, $\Delta R=1.50$ mag, $\Delta V=1.57$ mag, and $\Delta B=2.20$ mag.

It can be seen from Fig. 3 that the star spends most of the time at high light. During our photometric monitoring, several fading events of V521 Cyg are registered. Ibryamov et al. (2015a) described seven deep declines in the brightness observed in all bands. The drops in the brightness are non-periodic and have different duration and amplitudes. It is possible to suggest the existence of a frequent deep fading events during periods with insufficient data. The new data shows one fading event observed in the beginning of 2015 with

⁴ The tables with photometric CCD observations and data of the stars from our study only available at the Online material of the present paper.




Fig. 3. CCD *IRVB* light curves of V521 Cyg for the period 1997 June–2015 September

amplitudes 0.40 mag in *I*-band, 0.48 mag in *R*-band, 0.57 mag in *V*-band, and 0.81 mag for *B*-band.

The results obtained during our long-term study support the hypothesis of variable extinction as a reason for variability of V521 Cyg (Mitskevich & Pavlenko (2001) and Grankin et al. (2007)). The large amplitudes of observed declines in the brightness of the star (Fig. 3) are indication of UXor-type variability and the deep fading events presumably are result of circumstellar dust or clouds obscuration.

The measured colour indices $V - R$, $V - R$, and $B - V$ versus the stellar *V* magnitude during the period of our observations are plotted in Fig. 4. Nevertheless of the large amplitude of variability, the substantial bluing effect is not seen on the figure. A possible explanation of this phenomenon is that the declines in the brightness at different periods are caused by clouds of dust and particles of various sizes. It can be supposed that in this case we observe eclipses from a combination of clouds with different structure rotating in orbits around the star.

The *BVRI* light curves of V521 Cyg look very similar to light curves of FHO 27, which was classified as UXor by Ibryamov (2015b). FHO 27 is located at about $9'$ from V521 Cyg.




Fig. 4. Relationship between V magnitude and the $V - I$, $V - R$, and $B - V$ colour indices of V521 Cyg in the period 1997–2015

3.2. V752 Cyg

The variability of V752 Cyg was reported in Erastova & Tsvetkov (1978). Kohoutek & Wehmeyer (1997) confirmed the variability of the star, measured its photographic magnitude $m_{pg}=15.30$ mag and registered $H\alpha$ emission line in its spectrum. V752 Cyg is included in the list of YSO candidates published in Guieu et al. (2009).

The $BVRI$ light curves of V752 Cyg from all our CCD observations (Poljančič Beljan et al. 2014; Ibryamov 2015a; and the present paper) are shown in Fig. 5. The symbols used for the different telescopes are as in Fig. 3. The results from our recent multicolour CCD observations of V752 Cyg are summarized in Table 5. The columns have the same contents as in Table 4.

It can be seen from Fig. 5 that during our study the brightness of V752 Cyg vary around some intermediate level. The star shows principally increases in the brightness with different amplitudes. The brightness of the star during the period of our observations 2006–2015 vary in the range 14.28–15.55 mag for I -band, 14.89–16.67 mag for R -band, 15.38–17.71 mag for V -band, and 16.02–18.89 mag in B -band. The observed amplitudes are $\Delta I=1.27$ mag, $\Delta R=1.78$ mag, $\Delta V=2.33$ mag, and $\Delta B=2.87$ mag.

The colour indices $V - I$, $V - R$, and $V - B$ versus V magnitude during the period of our observations are plotted in Fig. 6. From the figure, it is seen that the star becomes bluer when it increases its brightness. Such colour variations are indication for flare events. These flares can be explained by increase in accretion rate from the circumstellar disk onto the stellar surface. Evidences of




Fig. 5. CCD *IRVB* light curves of V752 Cyg for the period 2006 December–2015 September

periodicity in the brightness variability are not detected. In the $J - H/H - K_s$ diagram (Fig. 2) V752 Cyg lies by 0.05 mag above the intrinsic T Tauri line – it have negligible infrared excess.

3.3. V1538 Cyg

The star V1538 Cyg was discovered and classified as a flare star by Erastova & Tsvetkov (1974). The authors reported a flare event on 1973 July 30 and determined its photographic magnitude $m_{pg}=18.50$ mag and $\Delta U=2.50$ mag. Laugalys et al. (2006) registered $H\alpha$ emission-line in the star’s spectrum and measured its magnitude $V=17.01$ mag and colour index $U - V=4.32$ mag. The authors determined the spectral type as M2 and the distance to the star as $r=248$ pc.

The spectrum of V1538 Cyg taken on 2007 October 23 by Corbally et al. (2009) shows a M1 photosphere with low emission in $H\alpha$. In $J - H/H - K_s$ diagram the star lies in the upper part of the early M dwarf band, so there are no indications of the presence of the envelope. This is confirmed by the *Spitzer* fluxes – the energy distribution of the star no showed infrared radiation. According to Corbally et al. (2009) V1538 Cyg can be either an unreddened field dwarf with chromospheric activity, located at 350 pc, or a post-T Tauri star at the front edge of the NGC 7000/IC 5070 complex. Armond et al. (2011) included V1538 Cyg in the list of $H\alpha$ emission-line star and measured its magnitudes as $I=16.28$, $R=16.85$ and $V=17.82$ mag.




Fig. 6. Relationship between V magnitude and the $V - I$, $V - R$, and $B - V$ colour indices of V752 Cyg in the period 2006–2015

The multicolour $BVRI$ light curves of V1538 Cyg from all our CCD observations in the period 2000–2015 (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 7. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 6. The columns have the same contents as in Table 4.

From Fig. 7 can be seen that during the period of our observations V1538 Cyg shows irregular variability in all bands, but significant flare events are not registered. The brightness of the star during the period 2000–2015 vary in the range 14.79–15.13 mag for I -band, 15.78–16.26 mag for R -band, 16.80–17.21 mag for V -band, and 18.11–18.64 mag for B -band. The observed amplitudes are $\Delta I=0.34$ mag, $\Delta R=0.48$ mag, $\Delta V=0.41$ mag, and $\Delta B=0.53$ mag in the same period.

Usually, such low amplitude variability is typical for low-mass WTTS, whose variability is produced by rotation of the spotted surface. This is confirmed by the position of the star in the 2MASS two-colour diagram in Fig. 2 – there are no indications of the presence of infrared excess. Evidences of periodicity in the brightness variability of V1538 Cyg are not detected.

3.4. V1539 Cyg

The star V1539 Cyg was discovered and classified as a $H\alpha$ emission-line star by Herbig (1958) with photographic magnitude $m_{pg}=17.00$ mag. Welin (1973)




Fig. 7. CCD *IRVB* light curves of V1538 Cyg for the period 2000 June–2015 September

confirmed it as a $H\alpha$ emission star with strong $H\alpha$ intensity and determined $V=15.50$ and $B=17.00$ mag.

Giesekeing & Schumann (1976) discovered that star’s brightness vary with amplitude greater than 0.25 mag, which led to the conclusion of the suspected variability. Herbig & Bell (1988) classified V1539 Cyg as a T Tauri star with $m_{pg}=16.30$ mag. Kohoutek & Wehmeyer (1997) confirmed the variability of the star with maximum brightness in photoelectric *V* system 14.50 mag, and determined its spectral type as K6IV.

The spectrum of V1539 Cyg taken on 2007 October 21 by Corbally et al. (2009) shows a G5e photosphere with strong emission in $H\alpha$. CaII and OI lines are also registered in emission. In $J - H/H - K_s$ diagram the star lies by 0.12 mag above the intrinsic T Tauri line. The *Spitzer* fluxes exhibit the presence of a considerable thermal emission from the dust envelope (Corbally et al. 2009). According to Findeisen et al. (2013) V1539 Cyg shows bursts events and they reported average stellar magnitude $\overline{R}=14.60$ mag and $\Delta R=0.70$ mag.

The *BVRI* light curves of V1539 Cyg from all our CCD observations (Poljančič Beljan et al. 2014; Ibryamov 2015a; and the present paper) are shown in Fig. 8. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 7. The columns have the same contents as in Table 4.

The brightness of V1539 Cyg during the period of our observations 1997–2015 vary in the range 13.34–13.68 mag for *I*-band, 14.28–14.72 mag for *R*-band, 15.28–15.79 mag for *V*-band, and 16.56–17.26 mag for *B*-band. The observed amplitudes are $\Delta I=0.34$ mag, $\Delta R=0.44$ mag, $\Delta V=0.51$ mag, and $\Delta B=0.70$ mag. From Fig. 8 it can be seen that during the period of our ob-




Fig. 8. CCD *IRVB* light curves of V1539 Cyg for the period 1997 June–2015 September

servations V1539 Cyg shows variability in all bands. This variability include short rises and decreases of the star’s brightness with small amplitudes.

Important result from our long-term photometric study of V1539 Cyg is that during the whole period of observations the total star’s brightness gradually decreases. Using a linear approximation for all our data of the star, we calculated the following values for the rates of decreases: $0.0117 \text{ mag yr}^{-1}$ for *I*-band, $0.0119 \text{ mag yr}^{-1}$ for *R*-band, $0.0109 \text{ mag yr}^{-1}$ for *V*-band, and $0.0090 \text{ mag yr}^{-1}$ for *B*-band.

The light curves of the star gives grounds to predict different reasons for observed variability of the star – existence of hot and cool spots on the stellar surface, and/or irregular obscuration of the star by circumstellar material. Evidences of periodicity in the brightness variability are not detected.

3.5. V1716 Cyg

The variability of V1716 Cyg was discovered by Erastova & Tsvetkov (1978). The authors reported irregular brightness fluctuations and measured its photographic magnitudes $m_{pg}=17.00\text{--}17.50 \text{ mag}$ (limit) and $U=16.10\text{--}17.50 \text{ mag}$ (limit). Armond et al. (2011) included V1716 Cyg in the list of $H\alpha$ emission-line star and measured its magnitudes as $I=15.95$, $R=16.74$ and $V=17.93 \text{ mag}$. Findeisen et al. (2013) discovered two bursts events of the star, which are separated by 35 days, as the first one lasting 5–20 days, and the second one – 3 days. The authors reported average stellar magnitude $\overline{R}=16.50 \text{ mag}$ and $\Delta R=1.10 \text{ mag}$.

Poljančič Beljan et al. (2014) determined the period of V1716 Cyg, which is found to be 4.15 days. This periodicity is probably connected with the stellar rotation and the position of the dark spots on the stellar surface.

The *BVRI* light curves of V1716 Cyg from all our CCD observations (Poljančič Beljan et al. 2014; Ibryamov 2015a; and the present paper) are shown in Fig. 9. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 8. The columns have the same contents as in Table 4.




Fig. 9. CCD *IRVB* light curves of V1716 Cyg for the period 1997 June–2015 September

The brightness of V1716 Cyg during the period of our observations 1997–2015 vary in the range 14.43–15.36 mag for *I*-band, 15.28–16.68 mag for *R*-band, 16.14–17.93 mag for *V*-band, and 17.05–19.50 mag for *B*-band. The observed amplitudes are $\Delta I=0.93$ mag, $\Delta R=1.40$ mag, $\Delta V=1.79$ mag, and $\Delta B=2.45$ mag in the same period.

Ibryamov et al. (2015a) described four eruptive events of V1716 Cyg observed in all bands. The irregular flares can be explained with short-lived accretion-related events at the stellar surface or as flares from UV Cet-type. Other irregular variations of the star’s brightness, observed in Fig. 9 are with smaller amplitudes and are caused probably by rotating hot and cool spots on the stellar surface. These results are an indication that V1716 Cyg is likely CTTS.

In the $J - H/H - K_s$ diagram (Fig. 2) the star lies by 0.11 mag above the intrinsic T Tauri line, i.e., it have infrared excess and this is indication of the presence of the envelope. Our new data confirm the periodicity of V1716 Cyg found by Poljančič Beljan et al. (2014).

3.6. V1957 Cyg

V1957 Cyg was discovered and classified as a flare star by Chavushian & Jankovics (1985). The authors reported a flare event on 1979 October 19 when the brightness of the star in U -band increased from 18.00 to 16.00 mag. In the following series of observations performed in 1981 and 1984 flare events are not registered. Laugalys et al. (2006) classified V1957 Cyg as a K6 dwarf with possible $H\alpha$ emission. The authors measured $V=16.30$ mag and colour index $U - V=4.05$ mag, and determined the distance to the star as $r=481$ pc.

The spectrum of V1957 Cyg taken on 2007 October 22 by Corbally et al. (2009) shows a Ge photosphere with weak emission in $H\alpha$. The authors reported that in $J - H/H - K_s$ diagram the star lies close to the intrinsic position of M0V star, but it can be a reddened G-type star, too. IRAC magnitudes measured by *Spitzer* show no infrared excess. According to authors V1957 Cyg may have been in a particularly active phase at the time of photometric observations, and in quiet stage during the spectral observations.

The $BVRI$ light curves of the star from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 10. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 9. The columns have the same contents as in Table 4.




Fig. 10. CCD $IRVB$ light curves of V1957 Cyg for the period 2000 June–2015 September

The brightness of V1957 Cyg during the period 2000–2015 vary in the range 14.38–14.79 mag. for I -band, 15.20–15.70 mag. for R -band, 16.06–16.54 mag. for V -band, and 17.25–17.83 mag. for B -band. The observed amplitudes are $\Delta I=0.41$ mag, $\Delta R=0.50$ mag, $\Delta V=0.48$ mag, and $\Delta B=0.58$ mag during the same period.

It can be seen from Fig. 10 that the star's brightness increasing smoothly to the mid-2010, then it begins decrease smoothly to the mid-2014. From the mid-2014 so far the star's brightness vary around some intermediate level with very small amplitudes. From the observed amplitudes of V1957 Cyg and its position in the two-colour diagram (Figure 2) it can be assumed that the star is probably M dwarf with chromospheric activity. Evidences of periodicity in the brightness variability of the star are not detected.

3.7. V2051 Cyg

V2051 Cyg was discovered and classified as a flare star by Parsamian et al. (1994). The authors reported a flare event on 1977 September 07 when the brightness of the star in U -band increased from the plate limit to 14.00 mag with amplitude $\Delta m_U > 4.00$ mag.

Laugalys et al. (2006) classified V2051 Cyg as a possible T Tauri star with $V=16.59$ mag. The spectrum of the star taken on 2007 October 21 by Corbally et al. (2009) shows a M3.5e photosphere with weak emission in $H\alpha$. In $J-H/H-K_s$ diagram the star lies close to the intrinsic position of M3V stars with no infrared excess. Its energy distribution, constructed from the 2MASS and *Spitzer* observations, also suggest for a normal star without envelope. According to Corbally et al. (2009) V2051 Cyg is most probably a M dwarf with weak chromospheric activity and the distance to the star is $r=137$ pc, i.e., the star cannot have any relation to the NGC 7000/IC 5070 complex.

Poljančič Beljan et al. (2014) determined the period of the star, which is found to be 384 days, likely caused by precession of the circumstellar disk or by clouds of gas and dust orbiting the star.

The multicolour $BVRI$ light curves of V2051 Cyg from all our CCD observations in the period 2000–2015 (Poljančič Beljan et al. 2014; Ibryamov 2015a; and the present paper) are shown in Fig 11. The symbols used for the different telescopes are as in Fig 3. The results of our recent multicolour CCD observations of the star are summarized in Table 10. The columns have the same contents as in Table 4.

During the period of our observations flare events are not registered, except a few low amplitude increases in brightness in V - and B -band. The brightness of the star during the period 2000–2015 vary in the range 13.84–14.02 mag for I -band, 15.40–15.68 mag for R -band, 16.40–16.77 mag for V -band, and 17.80–18.35 mag for B -band. The observed amplitudes are $\Delta I=0.18$ mag, $\Delta R=0.28$ mag, $\Delta V=0.37$ mag, and $\Delta B=0.55$ mag in the same period. Low amplitude of variability and relatively stable light variations confirm the assumptions on the nature of V2051 Cyg as a M dwarf with weak chromospheric activity (Corbally et al. 2009).

3.8. LkH α 186

The star LkH α 186 was discovered and classified as a $H\alpha$ emission-line star by Herbig (1958) with photographic magnitude $m_{pg}=18.00$ mag. Herbig & Bell (1988) classified it as a T Tauri variable with spectral type K5. Cohen & Kuhi (1979) measured $V=18.00$ mag of the star. In the paper of Weintraub (1990) LkH α 186 is also classified as a T Tauri variable.




Fig. 11. CCD *IRVB* light curves of V2051 Cyg for the period 2000 June–2015 September

Guieu et al. (2009) included the star in their list of YSO candidates and measured $I=14.747$, $V=17.395$, and $B=19.322$ mag. Armond et al. (2011) included it in the list of $H\alpha$ emission-line star and measured $I=15.85$, $R=16.68$ and $V=18.12$ mag and classified it as a CTTS.

The *BVRI* light curves of LkH α 186 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 12. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 11. The columns have the same contents as in Table 4.

The brightness of the star during the period 1997–2015 vary in the range 14.39–14.82 mag for *I*-band, 15.65–16.42 mag for *R*-band, 16.91–17.68 mag for *V*-band, and 18.37–19.50 mag for *B*-band. The observed amplitudes are $\Delta I=0.43$ mag, $\Delta R=0.77$ mag, $\Delta V=0.77$ mag, and $\Delta B=1.13$ mag.

It can be seen from Fig. 12 that the star usually spends most of the time at low light. LkH α 186 exhibit increases of the brightness with different amplitudes probably due to the presence of hot spots on the stellar surface. The variability is produced by variable accretion from the circumstellar disk and it is typical of CTTS.

In the two-colour diagram (Fig. 2) the star lies by 0.37 mag above the intrinsic T Tauri line. The star have clear infrared excess, indicating the presence of a circumstellar disk. Evidences of periodicity in the brightness variability are not detected.

3.9. LkH α 187

The star LkH α 187 was discovered and classified as a $H\alpha$ emission-line star by Herbig (1958) with photographic magnitude $m_{pg}=18.50$. Herbig & Bell (1988)




Fig. 12. CCD *IRVB* light curves of LkH α 186 for the period 1997 June–2015 September

classified it as a T Tauri variable with spectral type K3. Cohen & Kuhi (1979) measured $V=18.10$ mag of the star. In the work of Weintraub (1990) LkH α 187 is also classified as a T Tauri variable.

Guieu et al. (2009) included the star in their list of YSO candidates and measured $I=14.811$, $V=17.639$, and $B=19.476$ mag. Armond et al. (2011) included it in the list of H α emission-line star and measured $I=16.00$, $R=17.05$ and $V=18.56$ mag and classified it as a CTTS.

The *BVRI* light curves of LkH α 187 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 13. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 12. The columns have the same contents as in Table 4.

The brightness of the star during the period 1997–2015 vary in the range 14.46–15.02 mag for *I*-band, 15.68–16.71 mag for *R*-band, 17.02–18.15 mag for *V*-band, and 18.66–19.85 mag for *B*-band. The observed amplitudes are $\Delta I=0.56$ mag, $\Delta R=1.03$ mag, $\Delta V=1.13$ mag, and $\Delta B=1.19$ mag.

It can be seen from Fig. 13 that similar to LkH α 186, LkH α 187 usually spends most of the time at low light. The star exhibit increases of the brightness with different amplitudes. This variability is due likely to the presence of hot spots on the stellar surface, i.e., it is produced by strong variable accretion from the circumstellar disk.

LkH α 187 shows variability typical of CTTS. In the 2MASS two-colour diagram (Fig. 2) the star lies by 0.38 mag above the intrinsic T Tauri line. LkH α 187 have clear infrared excess, indicating the presence of a circumstellar disk. Evidences of periodicity in the brightness variability are not detected.




Fig. 13. CCD *IRVB* light curves of LkH α 187 for the period 1997 June–2015 September

3.10. LkH α 189

The star LkH α 189 was discovered and classified as a H α emission-line star by Herbig (1958) with photographic magnitude $m_{pg}=17.50$. Welin (1973) confirmed it as an H α emission-line star. Herbig & Bell (1988) classified it as a T Tauri variable with spectral type K6 and photographic magnitude 17.00 mag. Cohen & Kuhl (1979) measured $V=17.00$ mag of the star. In the paper of Weintraub (1990) LkH α 189 is also classified as a T Tauri variable.

Laugalys et al. (2006) classified LkH α 189 as a T Tauri star with $V=16.69$ mag and photographic spectral type K6. The position of the star in the two-colour $J-H/H-K_s$ diagram obtained by the Laugalys et al. (2006) and Meyer et al. (1997) suggest increased disk thickness and interstellar reddening. Guieu et al. (2009) included the star in their list of YSO candidates and measured $I=13.936$, $V=16.376$, and $B=18.175$ mag. Armond et al. (2011) included it in the list of H α emission-line star and measured $I=15.03$, $R=15.80$ and $V=17.10$ mag and classified it as a CTTS.

The *BVRI* light curves of LkH α 189 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 14. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 13. The columns have the same contents as in Table 4.

The brightness of LkH α 189 during the period 1997–2015 vary in the range 13.76–14.30 mag for *I*-band, 14.99–15.76 mag for *R*-band, 16.25–17.06 mag for *V*-band, and 17.69–18.75 mag for *B*-band. The observed amplitudes are $\Delta I=0.54$ mag, $\Delta R=0.77$ mag, $\Delta V=0.81$ mag, and $\Delta B=1.06$ mag in the same period.




Fig. 14. CCD *IRVB* light curves of LkH α 189 for the period 1997 June–2015 September

During our study the brightness of LkH α 189 vary around some intermediate level (Fig. 14). Both increases and drops in the brightness with different amplitudes are observed. The observed drops of the star’s brightness likely are caused by the existence of cool spots or groups of spots on the stellar surface or by irregular obscuration of the star by circumstellar material. The observed increases of the star’s brightness probably are caused by the existence of hot spots on the stellar surface.

We carried out a periodicity search using our data from 2010 May to 2015 September with PERSEA Version 2.6 (written by G. Maciejewski on the ANOVA technique, Schwarzenberg-Cherny (1996)) and with PERIOD04 (Lenz & Breger 2005) softwares. Our time-series analysis of the data indices a 2.450980 ± 0.029612 days period and led to the ephemeris:

$$JD(Max) = 2455337.925020 + 2.450980 * E. \quad (1)$$

False Alarm Probability estimation was done by randomly deleting about 10%-15% of the data for about 30 times and then retuning the period of determination. The period and starting age (T_0) determination remain stable, even with a subsample with 30% of the data removed. Fig. 15 exhibits the V-band folded light curve of LkH α 189 according to the ephemeris (1). Data obtained in *BRI*-bands show the same shape on periodicity. The found period is stable during time interval of several years and it is typical rotational period of young low-mass star. The periodicity could be caused by rotation modulation of the dark spots on the stellar surface.




Fig. 15. V-band folded curve of LkH α 189

3.11. LkH α 191

The star LkH α 191 was discovered and classified as a H α emission-line star by Herbig (1958) with photographic magnitude $m_{pg}=14.00$. Herbig & Bell (1988) classified it as a T Tauri variable with photographic magnitude 12.08 and spectral type K0. Terranegra et al. (1994) classified LkH α 191 as CTTS with $V=12.95$ mag and determined its photometric spectral type as K0V. Fernandez & Eiroa (1996) presented the light curve in V-band of the star for the period 1988 July–1991 August. The light curve show approximately constant brightness (with variations in V-band from 12.89 to 12.99 mag).

Laugalys et al. (2006) measured $V=13.06$ mag, colour index $U - V=3.506$ mag and photometric spectral type K6e. Authors noticed the absence of a strong H α emission and presence of a small dust emission suggested no significant deviation of the star from the main sequence. The position of LkH α 191 in two-colour $J - H/H - K_s$ diagram is slightly below intrinsic T Tauri line. According to Laugalys et al. (2006) LkH α 191 has probably lost its envelope during the past 50 years.

Grankin et al. (2007) conducted long-term study of LkH α 191 during the period 1986 July–1995 August. The authors reported that the star's brightness in V vary from 12.87 to 13.09 mag. Artemenko et al. (2012) determined the period of the star, which is found to be 2.08 days.

The *BVRI* light curves of LkH α 191 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 16. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 14. The columns have the same contents as in Table 4.

The brightness of LkH α 191 during the period 2000–2015 vary in the range 11.48–11.66 mag for *I*-band, 12.07–12.33 mag for *R*-band, 12.74–12.99 mag for *V*-band, and 13.87–14.23 mag for *B*-band. The observed amplitudes are $\Delta I=0.18$ mag, $\Delta R=0.26$ mag, $\Delta V=0.25$ mag, and $\Delta B=0.36$ mag.

It can be seen from Fig. 16 that LkH α 191 not shows some significant photometric activity. Our data agree with the hypothesis of an CTTS star, but LkH α 191 may also be post-T Tauri star. We carried out a periodicity




Fig. 16. CCD *IRVB* light curves of LkH α 191 for the period 2000 October–2015 September

search using our data, but we could not confirm the findings of Artemenko et al. (2012).

3.12. [KW97] 53-11

The star [KW97] 53-11 was mentioned in the catalog of Kohoutek & Wehmeyer (1997, 1999) with brightness in the photovisual system $m_{pv}=17.20$ mag. Cohen & Kuhi (1979) reported $V=17.70$ mag of the star and determined its spectral type as M2. Armond et al. (2011) report a lack of H α emission from this star.

The *RI* light curves of [KW97] 53-11 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 17. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 15. The columns have the same contents as in Table 4.

The brightness of the star during the period 2000–2015 vary in the range 15.81–16.44 mag for *I*-band and 18.32–19.04 mag for *R*-band. The observed amplitudes are $\Delta I=0.63$ mag and $\Delta R=0.72$ mag in the same period. The magnitudes of the star in *V* and *B*-band are below the limit of telescopes used.

It can be seen from Fig. 17 that during our study the brightness of [KW97] 53-11 vary around some intermediate level. The star shows both increases in the brightness with different amplitudes and short fading events in the brightness. Probably the observed fading events are caused by the existence of cool spots or group of spots, and the observed increases of the star’s brightness likely are caused by the existence of hot spots on the stellar surface. Evidences of periodicity in the brightness variability of [KW97] 53-11 are not detected.




Fig. 17. CCD *IRVB* light curves of [KW97] 53-11 for the period 1997 June–2015 September

3.13. [KW97] 53-23

The star [KW97] 53-23 was mentioned in the catalog of Kohoutek & Wehmeyer (1997, 1999) with only equatorial coordinates and finding charts available. Guieu et al. (2009) included the star in their list of YSO candidates and measured its magnitudes $I=15.093$, $V=18.445$, and $B=20.649$ mag.

The *BVRI* light curves of [KW97] 53-23 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 18. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 16. The columns have the same contents as in Table 4.

It can be seen from Fig. 18 that [KW97] 53-23 shows both increases and drops in the brightness with different amplitudes. The brightness of the star during the period 1997–2015 vary in the range 14.74–15.34 mag for *I*-band, 16.28–17.73 mag for *R*-band, 18.01–19.49 mag for *V*-band, and 19.92–20.49 mag for *B*-band. The observed amplitudes are $\Delta I=0.60$ mag, $\Delta R=1.45$ mag, $\Delta V=1.48$ mag, and $\Delta B=0.57$ mag.

The deepest drop in the brightness of [KW97] 53-23 is observed during 2014. Due to the limit of our photometric data during the minimal brightness some magnitudes in *V*- and especially in *B*-band are not registered. We have only some *B*-band photometric data of the star collected with the 2-m RCC and the 1.3-m RC telescopes.

The observed increases in the brightness of [KW97] 53-23 are caused probably by the existence of hot spots on the stellar surface, typical of CTTS. The observed drops in the brightness are caused probably by the existence of cool spots on the stellar surface or by irregular obscuration of the star by circumstellar material.




Fig. 18. CCD *IRVB* light curves of [KW97] 53-23 for the period 1997 June–2015 September

On Fig. 19, the colour indices $V - I$ and $V - R$ versus V magnitude during the period of our observations are plotted. The diagrams (especially $V/V - R$) shows evidences for bluing effect and probably the observed deepest drop in brightness are caused by obscuration of the star from circumstellar material.

In the two-colour diagram (Fig. 2) [KW97] 53-23 lies by 0.48 mag above the intrinsic T Tauri line. Therefore, the star have clear infrared excess, indicating the presence of a circumstellar disk. Evidences of periodicity in the brightness variability of the star are not detected.

3.14. [KW97] 53-36

The star [KW97] 53-36 was mentioned in the catalog of Kohoutek & Wehmeyer (1997, 1999) with it's brightness in the photographic system $m_{pg}=12.70$ mag.

The *BVRI* light curves of [KW97] 53-36 from all our CCD observations (Poljančič Beljan et al. 2014; and the present paper) are shown in Fig. 20. The symbols used for the different telescopes are as in Fig. 3. The results of our recent multicolour CCD observations of the star are summarized in Table 17. The columns have the same contents as in Table 4.

It can be seen from Fig. 20 that [KW97] 53-36 not shows some significant photometric activity. The brightness of the star during the period 2006–2015 vary in the range 11.53–11.67 mag for *I*-band, 11.84–11.96 mag for *R*-band, 12.17–12.38 mag for *V*-band, and 12.98–13.17 mag for *B*-band. The observed amplitudes are $\Delta I=0.14$ mag, $\Delta R=0.12$ mag, $\Delta V=0.21$ mag, and $\Delta B=0.19$ mag in the same period.

In the two-colour diagram (Fig. 2) the star lies close to the intrinsic position of G0 stars with no infrared excess, i.e., there are no indications of the




Fig. 19. Relationship between V magnitude and the $V - I$ and $V - R$ colour indices of [KW97] 53-23 in the period 1997–2015




Fig. 20. CCD $IRVB$ light curves of [KW97] 53-36 for the period 2006 December–2015 September

presence of a circumstellar disk. Most probably the star is a G giant with weak chromospheric activity. Evidences of periodicity in the brightness variability are not detected.

4. Conclusion

We presented the light curves of 14 PMS stars located in NGC 7000/IC 5070 star-forming complex. Our results complement the previously rare insights into their photometry. Seven of studied stars (V752 Cyg, V1539 Cyg, V1716 Cyg, LkH α 186, LkH α 187, LkH α 191, [KW97] 53-23) shows characteristics for CTTS, one star (V1538 Cyg) probably is WTTS, for three objects (V1539 Cyg, LkH α 189, [KW97] 53-11) we waver between CTTS and WTTS, for their exact classification are necessary spectral observations, three stars (V1957 Cyg, V2051 Cyg, [KW97] 53-36) likely are evolved PMS stars, and one stars (V521 Cyg) from our study we found indications of UXor variability. For one investigated star, LkH α 189, we found periodicity – 2.45-day period.

We are continuing to collect photometric observations of the field of 'Gulf of Mexico'. Further regular observations of the PMS stars in the same field will be of a great importance for their exact classification and are encouraging.

Acknowledgements

This research has made use of the NASA's Astrophysics Data System Abstract Service, the SIMBAD database and the VizieR catalogue access tool, operated at CDS, Strasbourg, France. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. The authors thank the Director of Skinakas Observatory Prof. I. Papamastorakis and Prof. I. Papadakis for the award of telescope time.

References

- Armond, T., Reipurth, B., Bally, J., Aspin, C., 2011, *A&A*, 528, A125
- Artemenko, S. A., Grankin, K. N., Petrov, P., 2012, *Astronomy Letters*, 38, 783
- Bally, J., Ginsburg, A., Probst, R., Reipurth, B., Shirley, Y. L., Stringfellow, G. S., 2014, *AJ*, 148, 120
- Bertout, C., 1989, *ARA&A*, 27, 3515
- Bessell, M. S., Brett, J. M., 1988, *PASP*, 100, 1134
- Bibo, E. A., Thé, P. S., 1990, *A&A*, 236, 155
- Carpenter, J. M., 2001, *AJ*, 121, 2851
- Chavarría-K., C., Terranegra, L., Alcalá, J. M., Neri, L., 1989, *RMxAA*, 18, 178
- Chavushian, H. S., Jankovics, I., 1985, *IBVS*, 2814, 1
- Cohen, M., Kuhl, L. V., 1979, *ApJ Suppl.*, 41, 743
- Corbally, C. J., Straižys, V., Laugalys, V., 2009, *Balt. Astron.*, 18, 111
- Dullemond, C. P., van den Ancker, M. E., Acke, B., van Boekel, R., 2003, *ApJ*, 594, L47
- Erastova, L. K., Tsvetkov, M. K., 1974, *IBVS*, 909, 1
- Erastova, L. K., Tsvetkov, M. K., 1978, *Perem. Zvezdy, Byull.*, 21, 79
- Fernandez, M., Eiroa, C., 1996, *A&A*, 310, 143
- Filin, A. Ya., 1974, *Perem. Zvezdy, Prolozh.*, 2, 63

- Findeisen, K., Hillenbrand, L., Ofek, E., Levitan, D., Sesar, B., Laher, R., Surace, J., 2013, *ApJ*, 768, 93
- Giesecking, F., Schumann, J. D., 1976, *A&AS*, 26, 367
- Grankin, K. N., Melnikov, S. Yu., Bouvier, J., Herbst, W., Shevchenko, V. S., 2007, *A&AS*, 461, 183
- Grinin, V. P., Kiselev, N. N., Minikulov, N. Kh., Chernova, G. P., Voshchinnikov, N. V., 1991, *Ap&SS*, 186, 283
- Guieu, S., Rebull, L. M., Stauffer, J. R., Hillenbrand, L. A., Carpenter, J. M. et al., 2009, *ApJ*, 697, 787
- Herbig, G. H., 1958, *ApJ*, 128, 259
- Herbig, G. H., Bell, K. R., 1988, *Third catalog of emission-line stars of the Orion population, Lick Observatory Bulletin 1111, Santa Cruz: Lick Observatory*, 90
- Herbst, W., Shevchenko, V. S., 1999, *AJ*, 118, 1043
- Herbst, W., Eisloffel, J., Mundt, R., Scholz, A., 2007, *Protostars and Planets V, Ed. B. Reipurth, D. Jewitt and K. Keil*, 297
- Hoffmeister, C., 1949, *AN*, 278, 24
- Ibryamov, S. I., Semkov, E. H., Peneva, S. P., 2015a, *Bulgarian Astronomical Journal*, 22, 3
- Ibryamov, S. I., Semkov, E. H., Peneva, S. P., 2015b, *PASA*, 32, e021
- Joy, A. H., 1945, *ApJ*, 102, 168
- Kohoutek, L., Wehmeyer, R., 1997, *Abhandlungen aus der Hamburger Sternwarte, Band 11, Teil 1, 2*
- Kohoutek, L., Wehmeyer, R., 1999, *A&AS*, 134, 255
- Laugalys, V., Straizys, V., 2002, *Balt. Astr.*, 11, 205
- Laugalys, V., Straizys, V., Vrba, F. J., Boyle, R. P., Philip, A. G. D., Kazlauskas, A., 2006, *Balt. Astron.*, 15, 483
- Lenz, P., Breger, M., 2005, *CoAst*, 146, 53
- Ménard, F., Bertout, C., 1999, *The Origin of Stars and Planetary Systems, Ed. V. J. Lada and N. D. Kylafis*, 341
- Meyer, M. R., Calvet, N., Hillenbrand, L. A., 1997, *AJ*, 114, 288
- Miller, A. A., Hillenbrand, L. A., Covey, K. R., Poznanski, D., Silverman, J. M. et al., 2011, *ApJ*, 730, 80
- Mitskevich, M. P., Pavlenko, E. P., 2001, *Ap*, 44, 411
- Parsamian, E. S., Chavira, E., Gonzales, G., 1994, *IBVS*, 4046, 1
- Poljančič Beljan, I., Jurdana-Šepić, R., Semkov, E., Ibryamov, S., Peneva, S., Tsvetkov, M., 2014, *A&A*, 568, A49
- Samus, N. N., Durlevich, O. V. et al., 2009, *VizieR Online Data Catalog: General Catalogue of Variable Stars*, 1, 2025
- Schwarzenberg-Cherny, A., 1996, *ApJ*, 460, 107
- Semkov, E. H., Peneva, S. P., Munari, U., Milani, A., Valisa, P., 2010, *A&A*, 523, L3
- Semkov, E. H., Peneva, S. P., Munari, U., Tsvetkov, M. K., Jurdana-Šepić, R., de Miguel, E., Schwartz, R. D., Dimitrov, D. P., Kjurkchieva, D. P., Radeva, V. S., 2012, *A&A*, 542, A43
- Semkov, E. H., Peneva, S. P., Ibryamov, S. I., Dimitrov, D. P., 2014, *Bulgarian Astronomical Journal*, 20, 59
- Skrutskie, M. F., Cutri, R. M., Stiening, R., Weinberg, M. D., Schneider, S. et al., 2006, *AJ*, 131, 1163
- Straizys, V., Corbally, C. J., Laugalys, V., 2008, *Balt. Astron.*, 17, 125
- Terranegra, L., Chavarría-K., C., Diaz, S., Gonzales-Patino, D., 1994, *A&AS*, 104, 557
- Voshchinnikov, N. V., 1989, *Astrofizika*, 30, 509
- Weintraub, D. A., 1990, *ApJS*, 74, 575
- Welin, G., 1973, *A&AS*, 9, 183

A ONLINE MATERIAL. Photometric CCD observations and data of the stars from our study

Table 4: Photometric CCD observations and data of V521 Cyg during the period 2014 August–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
03.08.2014	56873.386	12.25	13.13	14.02	15.35	Sch	FLI
04.08.2014	56874.407	12.27	13.11	13.97	15.33	Sch	FLI
18.08.2014	56888.450	12.09	12.93	13.77	14.99	Sch	FLI
19.08.2014	56889.365	12.04	12.85	13.68	14.93	Sch	FLI
29.08.2014	56899.303	11.93	12.74	13.55	14.77	1.3-m	ANDOR
26.11.2014	56988.181	11.95	12.74	13.53	14.74	Sch	FLI
13.12.2014	57005.181	11.91	12.71	13.50	14.69	Sch	FLI
14.12.2014	57006.220	11.94	12.73	13.51	-	Sch	FLI
24.12.2014	57016.174	12.05	12.87	13.74	15.01	2-m	VA
25.12.2014	57017.175	12.08	12.90	13.75	15.03	2-m	VA
21.02.2015	57074.641	12.31	13.22	14.10	15.50	Sch	FLI
23.04.2015	57136.487	12.08	12.92	13.72	15.00	Sch	FLI
25.04.2015	57138.491	12.14	13.01	13.85	15.14	Sch	FLI
18.05.2015	57161.423	12.02	12.83	13.64	14.87	Sch	FLI
19.05.2015	57162.460	11.98	12.77	13.57	14.80	Sch	FLI
21.05.2015	57164.459	12.02	12.84	13.62	14.89	Sch	FLI
24.05.2015	57167.409	12.02	12.81	13.65	14.87	2-m	VA
12.06.2015	57186.444	11.98	12.79	13.56	14.81	Sch	FLI
13.06.2015	57187.478	12.06	12.85	13.68	14.90	2-m	VA
16.06.2015	57190.367	12.08	12.86	13.70	14.96	2-m	VA
16.07.2015	57220.364	12.09	12.93	13.75	15.01	Sch	FLI
17.07.2015	57221.417	12.05	12.88	13.70	14.96	Sch	FLI
19.07.2015	57223.400	12.09	12.87	13.77	14.99	2-m	VA
20.07.2015	57224.422	12.06	12.85	13.70	14.95	2-m	VA
11.08.2015	57246.355	11.88	12.66	13.46	14.65	1.3-m	ANDOR
12.08.2015	57247.394	11.84	12.63	13.41	14.60	1.3-m	ANDOR
17.08.2015	57252.384	11.89	12.65	13.45	14.65	2-m	VA
24.08.2015	57259.337	11.89	12.69	13.43	14.65	Sch	FLI
25.08.2015	57260.342	11.88	12.67	13.43	14.63	Sch	FLI
03.09.2015	57269.328	11.90	12.69	13.44	14.69	Sch	FLI
04.09.2015	57270.330	11.93	12.69	13.52	14.71	2-m	VA
05.09.2015	57271.323	11.94	12.69	13.51	14.70	2-m	VA
06.09.2015	57272.288	11.94	12.71	13.56	14.72	2-m	VA

Table 5: Photometric CCD observations and data of V752 Cyg during the period 2014 August–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
03.08.2014	56873.386	15.10	16.37	17.43	18.36	Sch	FLI
04.08.2014	56874.407	14.96	15.76	16.36	16.96	Sch	FLI
18.08.2014	56888.450	15.11	15.98	16.60	17.30	Sch	FLI
19.08.2014	56889.365	15.21	16.33	17.20	18.19	Sch	FLI
26.11.2014	56988.181	15.07	16.06	16.70	17.62	Sch	FLI
13.12.2014	57005.181	14.97	16.10	16.87	17.91	Sch	FLI
14.12.2014	57006.220	15.21	16.43	17.16	-	Sch	FLI
21.02.2015	57074.641	15.25	16.28	16.90	17.97	Sch	FLI

Table 5: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
23.04.2015	57136.487	15.21	16.60	17.47	18.77	Sch	FLI
19.05.2015	57162.460	15.29	16.57	17.27	18.75	Sch	FLI
12.06.2015	57186.444	14.96	15.89	16.50	17.39	Sch	FLI
16.07.2015	57220.364	14.88	15.84	16.60	17.47	Sch	FLI
17.07.2015	57221.417	14.97	15.96	16.68	17.50	Sch	FLI
24.08.2015	57259.337	14.93	15.78	16.22	17.06	Sch	FLI
25.08.2015	57260.342	15.30	16.30	16.91	17.95	Sch	FLI
03.09.2015	57269.328	15.03	16.11	16.71	17.79	Sch	FLI

Table 6: Photometric CCD observations and data of V1538 Cyg during the period 2013 May–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
15.05.2013	56428.404	15.04	15.93	17.04	18.14	60-cm	FLI
10.04.2013	56392.510	15.03	16.12	17.11	18.44	Sch	FLI
11.04.2013	56394.482	15.00	15.99	16.94	18.40	Sch	FLI
02.05.2013	56415.418	14.97	16.03	16.95	18.43	Sch	FLI
17.05.2013	56430.408	15.00	15.95	16.96	-	60-cm	FLI
19.05.2013	56432.405	14.98	15.99	16.81	-	60-cm	FLI
30.05.2013	56443.367	14.96	16.00	16.97	18.50	Sch	FLI
31.05.2013	56444.354	14.96	15.98	16.97	18.45	Sch	FLI
04.08.2013	56509.288	15.00	16.06	17.05	18.47	Sch	FLI
05.08.2013	56510.369	15.04	16.02	16.96	18.19	60-cm	FLI
05.08.2013	56510.386	14.97	16.05	17.00	18.35	Sch	FLI
06.08.2013	56511.411	14.99	16.04	17.00	18.41	Sch	FLI
06.08.2013	56511.413	15.05	16.11	16.95	18.11	60-cm	FLI
07.08.2013	56512.398	14.99	16.04	16.98	18.35	Sch	FLI
07.08.2013	56512.403	15.06	16.01	17.02	18.10	60-cm	FLI
08.08.2013	56513.382	15.02	16.06	16.99	18.37	60-cm	FLI
09.08.2013	56514.350	15.02	15.96	17.00	18.41	60-cm	FLI
12.08.2013	56517.284	15.01	15.98	16.72	18.54	60-cm	FLI
04.09.2013	56540.274	14.96	16.00	16.97	18.36	Sch	FLI
05.09.2013	56541.322	14.96	15.99	16.94	18.27	Sch	FLI
06.09.2013	56542.381	15.00	16.09	17.00	18.40	Sch	FLI
11.09.2013	56547.383	15.07	16.03	16.98	18.21	60-cm	FLI
14.09.2013	56550.363	15.00	15.97	16.94	18.15	60-cm	FLI
17.09.2013	56553.282	14.94	15.99	16.97	18.37	1.3-m	ANDOR
11.10.2013	56577.306	15.02	15.98	16.98	18.23	60-cm	FLI
12.10.2013	56578.331	15.08	16.03	17.03	18.44	60-cm	FLI
07.11.2013	56604.264	15.02	16.08	17.06	18.46	60-cm	FLI
28.12.2013	56655.200	14.98	16.04	16.99	18.31	Sch	FLI
29.12.2013	56656.180	14.95	16.04	17.00	18.31	Sch	FLI
30.12.2013	56657.193	15.01	16.05	16.98	18.33	Sch	FLI
23.01.2014	56681.189	14.96	16.01	17.01	18.31	Sch	FLI
22.03.2014	56738.590	14.98	16.05	16.97	18.33	Sch	FLI
21.05.2014	56799.425	14.97	15.99	16.99	18.49	Sch	FLI
28.06.2014	56837.417	14.99	16.05	17.02	18.51	Sch	FLI
29.06.2014	56838.398	14.99	16.07	17.07	18.36	Sch	FLI
20.07.2014	56859.390	14.97	15.97	16.95	18.46	60-cm	FLI
21.07.2014	56860.391	15.02	16.06	17.11	-	60-cm	FLI
24.07.2014	56863.339	15.03	16.11	-	18.40	Sch	FLI
25.07.2014	56864.356	15.02	15.94	-	-	Sch	FLI
03.08.2014	56873.386	14.99	16.11	17.09	18.51	Sch	FLI

Table 6: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
04.08.2014	56874.407	15.01	16.14	17.15	18.58	Sch	FLI
18.08.2014	56888.450	14.98	16.04	16.99	18.39	Sch	FLI
19.08.2014	56889.365	14.96	16.04	16.99	18.35	Sch	FLI
29.08.2014	56899.303	14.99	16.06	16.98	18.36	1.3-m	ANDOR
26.11.2014	56988.181	14.98	16.02	16.95	18.45	Sch	FLI
13.12.2014	57005.181	14.98	16.10	17.06	18.41	Sch	FLI
14.12.2014	57006.220	14.98	16.04	16.93	-	Sch	FLI
21.02.2015	57074.641	14.96	15.96	16.98	18.35	Sch	FLI
23.04.2015	57136.487	14.94	15.96	16.88	18.20	Sch	FLI
25.04.2015	57138.491	14.95	16.00	16.90	18.40	Sch	FLI
18.05.2015	57161.423	14.94	15.95	16.87	18.53	Sch	FLI
19.05.2015	57162.460	14.94	15.99	16.91	18.41	Sch	FLI
21.05.2015	57164.459	14.95	15.98	16.87	18.23	Sch	FLI
12.06.2015	57186.444	14.95	15.99	16.88	18.24	Sch	FLI
16.07.2015	57220.364	14.94	15.96	16.92	18.26	Sch	FLI
17.07.2015	57221.417	14.94	15.99	16.92	18.24	Sch	FLI
11.08.2015	57246.355	14.92	15.96	16.91	18.34	1.3-m	ANDOR
12.08.2015	57247.394	13.96	15.99	16.94	18.37	1.3-m	ANDOR
24.08.2015	57259.337	14.97	15.99	16.90	18.32	Sch	FLI
25.08.2015	57260.342	14.95	16.03	16.89	18.33	Sch	FLI
03.09.2015	57269.328	14.96	15.95	16.86	18.23	Sch	FLI

Table 7: Photometric CCD observations and data of V1539 Cyg during the period 2014 August–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
03.08.2014	56873.386	13.57	14.59	15.65	17.01	Sch	FLI
04.08.2014	56874.407	13.49	14.47	15.52	16.85	Sch	FLI
18.08.2014	56888.450	13.61	14.66	15.74	17.13	Sch	FLI
19.08.2014	56889.365	13.60	14.62	15.71	17.14	Sch	FLI
29.08.2014	56899.303	13.58	14.61	15.66	17.04	1.3-m	ANDOR
26.11.2014	56988.181	13.61	14.66	15.72	17.13	Sch	FLI
13.12.2014	57005.181	13.54	14.57	15.54	16.98	Sch	FLI
14.12.2014	57006.220	13.60	14.65	15.62	-	Sch	FLI
21.02.2015	57074.641	13.59	14.58	15.54	16.85	Sch	FLI
23.04.2015	57136.487	13.53	14.54	15.58	16.97	Sch	FLI
25.04.2015	57138.491	13.60	14.64	15.68	17.03	Sch	FLI
18.05.2015	57161.423	13.62	14.67	15.72	17.05	Sch	FLI
19.05.2015	57162.460	13.54	14.54	15.54	16.89	Sch	FLI
21.05.2015	57164.459	13.58	14.63	15.67	17.07	Sch	FLI
12.06.2015	57186.444	13.52	14.56	15.52	16.90	Sch	FLI
16.07.2015	57220.364	13.57	14.62	15.69	17.08	Sch	FLI
17.07.2015	57221.417	13.62	14.67	15.72	17.09	Sch	FLI
11.08.2015	57246.355	13.53	14.60	15.68	17.08	1.3-m	ANDOR
12.08.2015	57247.394	13.58	14.63	15.68	17.08	1.3-m	ANDOR
24.08.2015	57259.337	13.60	14.64	15.67	17.04	Sch	FLI
25.08.2015	57260.342	13.58	14.63	15.62	17.09	Sch	FLI
03.09.2015	57269.328	13.52	14.56	15.53	16.94	Sch	FLI

Table 8: Photometric CCD observations and data of V1716 Cyg during the period 2014 August–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
03.08.2014	56873.386	15.09	16.48	17.67	19.29	Sch	FLI
04.08.2014	56874.407	15.07	16.58	17.66	-	Sch	FLI
18.08.2014	56888.450	15.14	16.59	17.82	-	Sch	FLI
19.08.2014	56889.365	15.15	16.64	17.77	19.25	Sch	FLI
29.08.2014	56899.303	14.99	16.41	17.59	19.21	1.3-m	ANDOR
26.11.2014	56988.181	15.15	16.64	17.75	19.45	Sch	FLI
13.12.2014	57005.181	15.09	16.64	17.64	-	Sch	FLI
14.12.2014	57006.220	15.09	16.52	17.59	-	Sch	FLI
24.12.2014	57016.174	14.99	16.24	17.49	19.18	2-m	VA
25.12.2014	57017.175	15.10	16.42	17.67	19.35	2-m	VA
21.02.2015	57074.641	15.14	16.58	17.64	-	Sch	FLI
23.04.2015	57136.487	15.02	16.40	17.49	19.15	Sch	FLI
25.04.2015	57138.491	15.08	16.52	17.61	19.07	Sch	FLI
18.05.2015	57161.423	15.03	16.42	17.66	-	Sch	FLI
19.05.2015	57162.460	15.15	16.61	17.56	19.20	Sch	FLI
21.05.2015	57164.459	15.05	16.44	17.53	19.32	Sch	FLI
24.05.2015	57167.409	15.04	16.24	17.57	19.42	2-m	VA
12.06.2015	57186.444	15.04	16.50	17.61	19.32	Sch	FLI
13.06.2015	57187.478	15.03	16.24	17.59	19.22	2-m	VA
16.06.2015	57190.367	14.93	16.15	17.45	19.14	2-m	VA
16.07.2015	57220.364	15.16	16.54	17.68	19.26	Sch	FLI
17.07.2015	57221.417	15.08	16.50	17.63	19.15	Sch	FLI
19.07.2015	57223.400	14.98	16.21	17.51	19.17	2-m	VA
20.07.2015	57224.422	15.05	16.29	17.61	19.29	2-m	VA
11.08.2015	57246.355	15.06	16.47	17.63	19.24	1.3-m	ANDOR
12.08.2015	57247.394	15.01	16.42	17.57	19.14	1.3-m	ANDOR
17.08.2015	57252.384	14.99	16.21	17.50	19.16	2-m	VA
24.08.2015	57259.337	15.05	16.49	17.45	-	Sch	FLI
25.08.2015	57260.342	15.02	16.39	17.49	18.92	Sch	FLI
03.09.2015	57269.328	15.09	16.46	17.59	19.22	Sch	FLI
04.09.2015	57270.330	15.05	16.33	17.63	19.29	2-m	VA
05.09.2015	57271.323	15.06	16.26	17.54	19.22	2-m	VA
06.09.2015	57272.288	15.00	16.18	17.44	19.08	2-m	VA

Table 9: Photometric CCD observations and data of V1957 Cyg during the period 2013 May–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
15.05.2013	56428.404	14.65	15.41	16.31	17.52	60-cm	FLI
10.04.2013	56392.510	14.57	15.45	16.32	-	Sch	FLI
11.04.2013	56394.482	14.63	15.47	16.33	17.55	Sch	FLI
02.05.2013	56415.418	14.63	15.47	16.33	17.60	Sch	FLI
17.05.2013	56430.408	14.65	15.52	16.29	-	60-cm	FLI
19.05.2013	56432.405	14.62	15.40	16.26	17.39	60-cm	FLI
30.05.2013	56443.367	14.57	15.44	16.30	17.50	Sch	FLI
31.05.2013	56444.354	11.57	15.43	16.26	17.47	Sch	FLI
04.08.2013	56509.288	14.65	15.62	16.47	17.80	Sch	FLI
05.08.2013	56510.369	14.69	15.59	16.40	17.54	60-cm	FLI
05.08.2013	56510.386	14.64	15.55	16.41	17.67	Sch	FLI
06.08.2013	56511.411	14.59	15.49	16.32	17.62	Sch	FLI
06.08.2013	56511.413	14.66	15.41	16.33	17.50	60-cm	FLI
07.08.2013	56512.398	14.58	15.46	16.29	17.55	Sch	FLI

Table 9: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
07.08.2013	56512.403	14.65	15.41	16.25	17.56	60-cm	FLI
08.08.2013	56513.382	14.63	15.54	16.23	17.55	60-cm	FLI
09.08.2013	56514.350	14.70	15.59	16.40	17.52	60-cm	FLI
12.08.2013	56517.284	14.55	15.41	16.28	17.72	60-cm	FLI
04.09.2013	56540.274	14.63	15.54	16.39	17.66	Sch	FLI
05.09.2013	56541.322	14.68	15.63	16.48	17.71	Sch	FLI
06.09.2013	56542.381	14.64	15.57	16.44	17.66	Sch	FLI
11.09.2013	56547.383	14.72	15.59	16.47	17.69	60-cm	FLI
14.09.2013	56550.363	14.68	15.50	16.38	17.66	60-cm	FLI
17.09.2013	56553.282	14.58	15.51	16.36	17.56	1.3-m	ANDOR
11.10.2013	56577.306	14.66	15.58	16.37	17.58	60-cm	FLI
12.10.2013	56578.331	14.73	15.56	16.43	17.70	60-cm	FLI
07.11.2013	56604.264	14.79	15.64	16.51	17.82	60-cm	FLI
28.12.2013	56655.200	14.60	15.49	16.36	17.52	Sch	FLI
29.12.2013	56656.180	14.66	15.60	16.44	17.68	Sch	FLI
30.12.2013	56657.193	14.67	15.60	16.49	17.81	Sch	FLI
23.01.2014	56681.189	14.60	15.48	16.33	17.60	Sch	FLI
22.03.2014	56738.590	14.63	15.52	16.35	17.54	Sch	FLI
21.05.2014	56799.425	14.67	15.60	16.43	17.73	Sch	FLI
28.06.2014	56837.417	14.63	15.49	16.39	17.69	Sch	FLI
29.06.2014	56838.398	14.61	15.55	16.37	17.61	Sch	FLI
20.07.2014	56859.390	14.68	15.50	16.30	-	60-cm	FLI
21.07.2014	56860.391	14.70	15.56	16.32	-	60-cm	FLI
24.07.2014	56863.339	14.65	15.58	-	17.67	Sch	FLI
25.07.2014	56864.356	14.62	15.55	-	-	Sch	FLI
03.08.2014	56873.386	14.66	15.60	16.45	17.68	Sch	FLI
04.08.2014	56874.407	14.66	15.58	16.42	17.69	Sch	FLI
18.08.2014	56888.450	14.68	15.68	16.54	17.76	Sch	FLI
19.08.2014	56889.365	14.66	15.61	16.46	17.68	Sch	FLI
29.08.2014	56899.303	14.69	15.64	16.49	17.75	1.3-m	ANDOR
26.11.2014	56988.181	14.74	15.70	16.54	17.83	Sch	FLI
13.12.2014	57005.181	14.65	15.59	16.40	17.72	Sch	FLI
14.12.2014	57006.220	14.69	15.62	16.36	-	Sch	FLI
21.02.2015	57074.641	14.69	15.60	16.40	17.67	Sch	FLI
23.04.2015	57136.487	14.64	15.61	16.41	17.70	Sch	FLI
25.04.2015	57138.491	14.67	15.60	16.40	17.73	Sch	FLI
18.05.2015	57161.423	14.70	15.63	16.48	17.82	Sch	FLI
19.05.2015	57162.460	14.71	15.66	16.48	17.80	Sch	FLI
21.05.2015	57164.459	14.67	15.60	16.41	17.62	Sch	FLI
12.06.2015	57186.444	14.66	15.60	16.39	17.71	Sch	FLI
16.07.2015	57220.364	14.70	15.64	16.46	17.68	Sch	FLI
17.07.2015	57221.417	14.68	15.60	16.45	17.75	Sch	FLI
11.08.2015	57246.355	14.67	15.62	16.47	17.75	1.3-m	ANDOR
12.08.2015	57247.394	14.65	15.58	16.43	17.72	1.3-m	ANDOR
24.08.2015	57259.337	14.66	15.57	16.40	17.70	Sch	FLI
25.08.2015	57260.342	14.66	15.63	16.38	17.68	Sch	FLI
03.09.2015	57269.328	14.66	15.57	16.34	17.67	Sch	FLI

Table 10: Photometric CCD observations and data of V2051 Cyg during the period 2014 August–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
03.08.2014	56873.386	13.93	15.65	16.58	18.12	Sch	FLI

Table 10: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
04.08.2014	56874.407	13.92	15.59	16.69	18.08	Sch	FLI
18.08.2014	56888.450	13.94	15.58	16.60	18.10	Sch	FLI
19.08.2014	56889.365	13.93	15.57	16.62	18.15	Sch	FLI
29.08.2014	56899.303	13.92	15.52	16.58	18.11	1.3-m	ANDOR
26.11.2014	56988.181	13.93	15.57	16.59	18.12	Sch	FLI
13.12.2014	57005.181	13.94	15.55	16.59	18.09	Sch	FLI
14.12.2014	57006.220	13.93	15.53	16.48	-	Sch	FLI
21.02.2015	57074.641	13.92	15.56	16.61	18.05	Sch	FLI
23.04.2015	57136.487	13.92	15.58	16.54	18.05	Sch	FLI
25.04.2015	57138.491	13.93	15.57	16.58	18.06	Sch	FLI
18.05.2015	57161.423	13.94	15.54	16.62	-	Sch	FLI
19.05.2015	57162.460	13.88	15.57	16.54	18.16	Sch	FLI
21.05.2015	57164.459	13.92	15.52	16.53	18.10	Sch	FLI
12.06.2015	57186.444	13.93	15.56	16.49	18.10	Sch	FLI
16.07.2015	57220.364	13.92	15.56	16.46	18.14	Sch	FLI
17.07.2015	57221.417	13.92	15.58	16.61	18.14	Sch	FLI
11.08.2015	57246.355	13.90	15.53	16.60	18.15	1.3-m	ANDOR
12.08.2015	57247.394	13.90	15.52	16.60	18.15	1.3-m	ANDOR
24.08.2015	57259.337	13.93	15.56	16.56	17.89	Sch	FLI
25.08.2015	57260.342	13.93	15.55	16.52	18.03	Sch	FLI
03.09.2015	57269.328	13.92	15.56	16.55	18.10	Sch	FLI

Table 11: Photometric CCD observations and data of LkH α 186 during the period 2013 April–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
10.04.2013	56392.510	14.44	15.99	17.10	-	Sch	FLI
11.04.2013	56394.482	14.46	16.02	17.19	18.68	Sch	FLI
02.05.2013	56415.418	14.59	16.19	17.40	19.12	Sch	FLI
15.05.2013	56428.404	14.54	15.98	17.22	-	60-cm	FLI
17.05.2013	56430.408	14.54	16.07	17.26	-	60-cm	FLI
19.05.2013	56432.405	14.65	16.16	17.48	-	60-cm	FLI
30.05.2013	56443.367	14.58	16.19	17.38	19.20	Sch	FLI
31.05.2013	56444.354	14.55	16.18	17.45	-	Sch	FLI
04.07.2013	56478.363	14.64	15.91	17.29	18.83	2-m	VA
01.08.2013	56506.281	14.61	15.91	17.27	18.89	2-m	VA
02.08.2013	56507.272	14.63	15.91	17.25	18.90	2-m	VA
03.08.2013	56508.318	14.57	15.85	17.17	18.78	2-m	VA
04.08.2013	56509.288	14.52	16.06	17.38	18.87	Sch	FLI
05.08.2013	56510.369	14.59	16.03	17.31	18.85	60-cm	FLI
05.08.2013	56510.386	14.52	16.08	17.34	18.97	Sch	FLI
06.08.2013	56511.411	14.54	16.13	17.35	18.88	Sch	FLI
06.08.2013	56511.413	14.61	16.17	17.33	-	60-cm	FLI
07.08.2013	56512.398	14.57	16.16	17.43	19.01	Sch	FLI
07.08.2013	56512.403	14.66	16.13	17.45	-	60-cm	FLI
08.08.2013	56513.382	14.56	16.09	17.32	18.80	60-cm	FLI
09.08.2013	56514.350	14.54	16.00	17.26	-	60-cm	FLI
12.08.2013	56517.284	14.56	16.07	17.11	18.74	60-cm	FLI
04.09.2013	56540.274	14.60	16.19	17.45	19.18	Sch	FLI
05.09.2013	56541.322	14.58	16.12	17.33	18.94	Sch	FLI
06.09.2013	56542.381	14.58	16.17	17.41	18.98	Sch	FLI
07.09.2013	56543.396	14.70	15.98	17.36	18.98	2-m	VA
08.09.2013	56544.264	14.77	16.08	17.45	19.07	2-m	VA

Table 11: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
11.09.2013	56547.383	14.63	16.06	17.38	18.95	60-cm	FLI
14.09.2013	56550.363	14.61	16.11	17.42	-	60-cm	FLI
17.09.2013	56553.282	14.58	16.06	17.30	18.79	1.3-m	ANDOR
11.10.2013	56577.306					60-cm	FLI
12.10.2013	56578.331					60-cm	FLI
07.11.2013	56604.264	14.65	16.17	17.53	19.15	60-cm	FLI
09.12.2013	56636.185	14.57	15.88	17.27	18.96	2-m	VA
28.12.2013	56655.200	14.55	16.19	17.50	19.15	Sch	FLI
29.12.2013	56656.180	14.53	16.04	17.25	18.66	Sch	FLI
30.12.2013	56657.193	14.51	16.09	17.31	18.55	Sch	FLI
23.01.2014	56681.189	14.75	16.42	17.68	19.35	Sch	FLI
05.02.2014	56694.194	-	15.94	17.32	-	2-m	VA
22.03.2014	56738.590	14.56	16.07	17.43	18.89	Sch	FLI
21.05.2014	56799.425	14.54	16.18	17.35	18.88	Sch	FLI
23.05.2014	56801.436	14.57	15.83	17.16	18.71	2-m	VA
23.06.2014	56832.391	14.58	15.83	17.14	18.75	2-m	VA
25.06.2014	56834.357	14.52	15.78	17.04	18.64	2-m	VA
26.06.2014	56835.462	14.41	15.69	16.91	18.43	2-m	VA
28.06.2014	56837.417	14.42	15.94	17.10	18.49	Sch	FLI
29.06.2014	56838.398	14.45	15.97	17.19	18.63	Sch	FLI
20.07.2014	56859.390	14.52	15.96	17.15	-	60-cm	FLI
21.07.2014	56860.391	14.54	16.06	17.11	-	60-cm	FLI
24.07.2014	56863.339	14.56	16.18	-	18.80	Sch	FLI
25.07.2014	56864.356	14.52	16.24	-	-	Sch	FLI
03.08.2014	56873.386	14.43	15.96	17.16	18.43	Sch	FLI
04.08.2014	56874.407	14.50	16.11	17.34	-	Sch	FLI
18.08.2014	56888.450	14.46	15.98	17.15	18.49	Sch	FLI
19.08.2014	56889.365	14.49	16.02	17.21	18.69	Sch	FLI
29.08.2014	56899.303	14.65	16.15	17.45	19.16	1.3-m	ANDOR
26.11.2014	56988.181	14.47	16.00	17.09	18.55	Sch	FLI
13.12.2014	57005.181	14.67	16.30	17.55	19.11	Sch	FLI
14.12.2014	57006.220	14.57	16.16	17.32	-	Sch	FLI
24.12.2014	57016.174	14.64	15.98	17.38	19.03	2-m	VA
25.12.2014	57017.175	14.71	16.04	17.42	19.08	2-m	VA
21.02.2015	57074.641	14.60	16.15	17.39	19.01	Sch	FLI
23.04.2015	57136.487	14.54	16.06	17.27	18.71	Sch	FLI
25.04.2015	57138.491	14.61	16.14	17.36	19.02	Sch	FLI
18.05.2015	57161.423	14.50	15.99	17.18	-	Sch	FLI
19.05.2015	57162.460	14.50	16.01	17.22	18.68	Sch	FLI
21.05.2015	57164.459	14.54	16.02	17.29	18.96	Sch	FLI
24.05.2015	57167.409	14.55	15.80	17.16	18.92	2-m	VA
12.06.2015	57186.444	14.50	15.99	17.15	18.85	Sch	FLI
13.06.2015	57187.478	14.55	15.82	17.20	18.74	2-m	VA
16.06.2015	57190.367	14.42	15.69	16.99	18.48	2-m	VA
16.07.2015	57220.364	14.49	15.95	17.05	18.53	Sch	FLI
17.07.2015	57221.417	14.53	16.06	17.28	18.73	Sch	FLI
19.07.2015	57223.400	14.57	15.84	17.18	18.81	2-m	VA
20.07.2015	57224.422	14.50	15.73	17.10	18.71	2-m	VA
11.08.2015	57246.355	14.42	15.88	17.07	18.49	1.3-m	ANDOR
12.08.2015	57247.394	14.52	15.99	17.20	18.69	1.3-m	ANDOR
17.08.2015	57252.384	14.39	15.65	16.92	18.37	2-m	VA
24.08.2015	57259.337	14.48	16.01	17.06	18.76	Sch	FLI
25.08.2015	57260.342	14.48	15.98	17.08	18.79	Sch	FLI
03.09.2015	57269.328	14.56	16.07	17.25	18.81	Sch	FLI
04.09.2015	57270.330	14.49	15.73	17.07	18.61	2-m	VA
05.09.2015	57271.323	14.58	15.84	17.16	18.86	2-m	VA
06.09.2015	57272.288	14.57	15.83	17.16	18.84	2-m	VA

Table 12: Photometric CCD observations and data of LkH α 187 during the period 2013 April–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
10.04.2013	56392.510	14.80	16.43	17.81	-	Sch	FLI
11.04.2013	56394.482	14.76	16.33	17.71	19.23	Sch	FLI
02.05.2013	56415.418	14.73	16.30	17.71	19.33	Sch	FLI
15.05.2013	56428.404	14.63	16.14	17.43	-	60-cm	FLI
17.05.2013	56430.408	14.63	16.15	17.31	-	60-cm	FLI
19.05.2013	56432.405	14.74	16.30	17.40	-	60-cm	FLI
30.05.2013	56443.367	14.70	16.18	17.70	-	Sch	FLI
31.05.2013	56444.354	14.71	16.28	17.78	-	Sch	FLI
04.07.2013	56478.363	14.77	16.11	17.61	19.42	2-m	VA
01.08.2013	56506.281	14.64	15.95	17.40	19.08	2-m	VA
02.08.2013	56507.272	14.66	15.97	17.48	19.11	2-m	VA
03.08.2013	56508.318	14.59	15.87	17.26	18.94	2-m	VA
04.08.2013	56509.288	14.62	16.16	17.44	19.29	Sch	FLI
05.08.2013	56510.369	14.72	16.25	17.69	-	60-cm	FLI
05.08.2013	56510.386	14.70	16.23	17.56	19.13	Sch	FLI
06.08.2013	56511.411	14.70	16.30	17.61	19.22	Sch	FLI
06.08.2013	56511.413	14.68	16.21	17.59	19.45	60-cm	FLI
07.08.2013	56512.398	14.68	16.28	17.65	19.28	Sch	FLI
07.08.2013	56512.403	14.69	16.22	17.76	19.36	60-cm	FLI
08.08.2013	56513.382	14.67	16.22	17.76	19.10	60-cm	FLI
09.08.2013	56514.350	14.62	16.16	17.51	18.81	60-cm	FLI
12.08.2013	56517.284	14.71	16.01	17.85	-	60-cm	FLI
04.09.2013	56540.274	14.63	16.09	17.34	18.86	Sch	FLI
05.09.2013	56541.322	14.66	16.21	17.47	19.13	Sch	FLI
06.09.2013	56542.381	14.69	16.28	17.53	19.16	Sch	FLI
07.09.2013	56543.396	14.76	16.01	17.42	19.14	2-m	VA
08.09.2013	56544.264	14.81	16.11	17.54	19.30	2-m	VA
11.09.2013	56547.383	14.70	16.22	17.80	-	60-cm	FLI
14.09.2013	56550.363	14.68	16.18	17.57	19.02	60-cm	FLI
17.09.2013	56553.282	14.64	16.19	17.52	19.20	1.3-m	ANDOR
11.10.2013	56577.306	14.66	16.16	-	-	60-cm	FLI
12.10.2013	56578.331	14.64	16.16	17.45	19.27	60-cm	FLI
07.11.2013	56604.264	14.72	16.27	17.72	19.36	60-cm	FLI
09.12.2013	56636.185	14.72	16.07	17.54	19.32	2-m	VA
28.12.2013	56655.200	14.70	16.29	17.77	-	Sch	FLI
29.12.2013	56656.180	14.71	16.29	17.65	19.20	Sch	FLI
30.12.2013	56657.193	14.67	16.27	17.60	19.03	Sch	FLI
23.01.2014	56681.189	14.64	16.18	17.48	19.21	Sch	FLI
05.02.2014	56694.194	-	16.00	17.46	19.55	2-m	VA
22.03.2014	56738.590	14.73	16.29	17.71	19.14	Sch	FLI
21.05.2014	56799.425	14.65	16.30	17.80	-	Sch	FLI
23.05.2014	56801.436	14.57	15.90	17.43	19.30	2-m	VA
23.06.2014	56832.391	14.62	15.87	17.28	19.00	2-m	VA
25.06.2014	56834.357	14.66	15.99	17.49	19.28	2-m	VA
26.06.2014	56835.462	14.60	15.88	17.33	19.09	2-m	VA
28.06.2014	56837.417	14.63	16.19	17.53	19.27	Sch	FLI
29.06.2014	56838.398	14.65	16.23	17.54	19.31	Sch	FLI
20.07.2014	56859.390	14.72	16.31	-	-	60-cm	FLI
21.07.2014	56860.391	14.60	16.08	17.31	-	60-cm	FLI
24.07.2014	56863.339	14.70	16.26	-	-	Sch	FLI
25.07.2014	56864.356	14.65	16.25	-	-	Sch	FLI
03.08.2014	56873.386	14.70	16.28	17.74	19.29	Sch	FLI
04.08.2014	56874.407	14.65	16.24	17.72	-	Sch	FLI
18.08.2014	56888.450	14.67	16.23	17.53	19.08	Sch	FLI
19.08.2014	56889.365	14.62	16.16	17.47	18.99	Sch	FLI
29.08.2014	56899.303	14.60	16.16	17.51	19.21	1.3-m	ANDOR

Table 12: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
26.11.2014	56988.181	14.61	16.18	17.46	19.29	Sch	FLI
13.12.2014	57005.181	14.62	16.18	17.48	19.36	Sch	FLI
14.12.2014	57006.220	14.69	16.30	17.64	-	Sch	FLI
24.12.2014	57016.174	14.63	16.03	17.50	19.32	2-m	VA
25.12.2014	57017.175	14.58	15.93	17.36	19.15	2-m	VA
21.02.2015	57074.641	14.61	16.09	17.35	19.18	Sch	FLI
23.04.2015	57136.487	14.69	16.27	17.60	19.15	Sch	FLI
25.04.2015	57138.491	14.65	16.14	17.50	19.35	Sch	FLI
18.05.2015	57161.423	14.71	16.16	17.55	-	Sch	FLI
19.05.2015	57162.460	14.62	16.22	17.52	19.39	Sch	FLI
21.05.2015	57164.459	14.71	16.28	17.67	19.07	Sch	FLI
24.05.2015	57167.409	14.57	15.86	17.29	19.13	2-m	VA
12.06.2015	57186.444	14.54	16.08	17.46	19.06	Sch	FLI
13.06.2015	57187.478	14.54	15.83	17.31	18.96	2-m	VA
16.06.2015	57190.367	14.59	15.92	17.42	19.30	2-m	VA
16.07.2015	57220.364	14.55	16.06	17.42	19.08	Sch	FLI
17.07.2015	57221.417	14.64	16.20	17.62	19.13	Sch	FLI
19.07.2015	57223.400	14.57	15.87	17.30	19.06	2-m	VA
20.07.2015	57224.422	14.64	15.94	17.46	19.34	2-m	VA
11.08.2015	57246.355	14.57	16.09	17.42	19.05	1.3-m	ANDOR
12.08.2015	57247.394	14.60	16.18	17.52	19.20	1.3-m	ANDOR
17.08.2015	57252.384	14.63	15.96	17.41	19.32	2-m	VA
24.08.2015	57259.337	14.63	16.15	17.41	19.29	Sch	FLI
25.08.2015	57260.342	14.63	16.20	17.55	19.15	Sch	FLI
03.09.2015	57269.328	14.61	16.15	17.51	19.10	Sch	FLI
04.09.2015	57270.330	14.56	15.82	17.21	18.99	2-m	VA
05.09.2015	57271.323	14.65	15.92	17.43	19.15	2-m	VA
06.09.2015	57272.288	14.46	15.68	17.02	18.66	2-m	VA

Table 13: Photometric CCD observations and data of LkH α 189 during the period 2013 May–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
15.05.2013	56428.404	13.82	15.12	16.34	17.87	60-cm	FLI
10.04.2013	56392.510	13.93	15.23	16.52	-	Sch	FLI
11.04.2013	56394.482	13.89	15.24	16.45	18.01	Sch	FLI
02.05.2013	56415.418	13.94	15.25	16.48	17.98	Sch	FLI
17.05.2013	56430.408	13.90	15.23	16.48	-	60-cm	FLI
19.05.2013	56432.405	13.92	15.22	16.47	17.96	60-cm	FLI
30.05.2013	56443.367	13.85	15.16	16.34	18.16	Sch	FLI
31.05.2013	56444.354	13.82	15.10	16.28	17.90	Sch	FLI
04.07.2013	56478.363	13.96	15.09	16.37	18.03	2-m	VA
01.08.2013	56506.281	13.93	15.11	16.40	18.12	2-m	VA
02.08.2013	56507.272	13.90	15.03	16.30	18.04	2-m	VA
03.08.2013	56508.318	13.97	15.16	16.46	18.18	2-m	VA
04.08.2013	56509.288	13.93	15.26	16.53	18.07	Sch	FLI
05.08.2013	56510.369	14.00	15.35	16.58	18.28	60-cm	FLI
05.08.2013	56510.386	14.00	15.35	16.61	18.31	Sch	FLI
06.08.2013	56511.411	13.94	15.30	16.53	18.32	Sch	FLI
06.08.2013	56511.413	13.94	15.30	16.56	18.18	60-cm	FLI
07.08.2013	56512.398	13.98	15.33	16.58	18.16	Sch	FLI
07.08.2013	56512.403	13.98	15.28	16.60	18.30	60-cm	FLI
08.08.2013	56513.382	13.96	15.31	16.57	18.18	60-cm	FLI

Table 13: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
09.08.2013	56514.350	13.86	15.22	16.45	18.15	60-cm	FLI
12.08.2013	56517.284	13.83	15.09	16.32	17.89	60-cm	FLI
04.09.2013	56540.274	13.94	15.32	16.56	18.22	Sch	FLI
05.09.2013	56541.322	13.86	15.20	16.41	17.99	Sch	FLI
06.09.2013	56542.381	13.90	15.28	16.48	18.05	Sch	FLI
07.09.2013	56543.396	13.99	15.16	16.47	18.18	2-m	VA
08.09.2013	56544.264	14.08	15.26	16.53	18.20	2-m	VA
11.09.2013	56547.383	14.07	15.43	16.73	18.67	60-cm	FLI
14.09.2013	56550.363	13.98	15.28	16.61	18.51	60-cm	FLI
17.09.2013	56553.282	13.95	15.33	16.61	18.27	1.3-m	ANDOR
11.10.2013	56577.306	13.99	15.29	16.55	18.44	60-cm	FLI
12.10.2013	56578.331	13.92	15.27	16.48	18.31	60-cm	FLI
07.11.2013	56604.264	14.13	15.57	16.82	18.64	60-cm	FLI
09.12.2013	56636.185	13.96	15.16	16.47	18.18	2-m	VA
28.12.2013	56655.200	13.93	15.32	16.57	18.12	Sch	FLI
29.12.2013	56656.180	13.88	15.21	16.39	18.00	Sch	FLI
30.12.2013	56657.193	14.02	15.38	16.62	18.30	Sch	FLI
23.01.2014	56681.189	13.93	15.27	16.50	18.13	Sch	FLI
05.02.2014	56694.194	-	15.42	16.78	18.71	2-m	VA
22.03.2014	56738.590	13.94	15.31	16.54	18.26	Sch	FLI
21.05.2014	56799.425	14.09	15.47	16.74	-	Sch	FLI
23.05.2014	56801.436	13.95	15.13	16.45	18.22	2-m	VA
23.06.2014	56832.391	14.04	15.26	16.58	18.36	2-m	VA
25.06.2014	56834.357	13.94	15.12	16.42	18.13	2-m	VA
26.06.2014	56835.462	13.90	15.10	16.40	18.10	2-m	VA
28.06.2014	56837.417	14.04	15.47	16.73	18.51	Sch	FLI
29.06.2014	56838.398	14.15	15.61	16.85	18.44	Sch	FLI
20.07.2014	56859.390	13.94	15.30	16.45	18.06	60-cm	FLI
21.07.2014	56860.391	14.06	15.43	16.69	-	60-cm	FLI
24.07.2014	56863.339	13.98	15.34	-	18.25	Sch	FLI
25.07.2014	56864.356	13.94	15.29	-	-	Sch	FLI
03.08.2014	56873.386	14.14	15.53	16.81	18.45	Sch	FLI
04.08.2014	56874.407	13.99	15.37	16.71	-	Sch	FLI
18.08.2014	56888.450	14.02	15.48	16.70	18.38	Sch	FLI
19.08.2014	56889.365	13.99	15.42	16.69	18.30	Sch	FLI
29.08.2014	56899.303	13.91	15.26	16.50	18.12	1.3-m	ANDOR
26.11.2014	56988.181	14.18	15.64	16.87	18.59	Sch	FLI
13.12.2014	57005.181	14.14	15.60	16.81	18.75	Sch	FLI
14.12.2014	57006.220	13.97	15.38	16.60	-	Sch	FLI
24.12.2014	57016.174	13.86	15.07	16.35	18.02	2-m	VA
25.12.2014	57017.175	14.06	15.31	16.62	18.42	2-m	VA
21.02.2015	57074.641	13.94	15.32	16.49	18.18	Sch	FLI
23.04.2015	57136.487	13.98	15.39	16.58	18.25	Sch	FLI
25.04.2015	57138.491	13.93	15.29	16.50	18.10	Sch	FLI
18.05.2015	57161.423	13.95	15.29	16.52	17.99	Sch	FLI
19.05.2015	57162.460	14.00	15.39	16.60	18.43	Sch	FLI
21.05.2015	57164.459	14.04	15.41	16.65	18.39	Sch	FLI
24.05.2015	57167.409	14.00	15.17	16.53	18.40	2-m	VA
12.06.2015	57186.444	14.05	15.44	16.66	18.39	Sch	FLI
13.06.2015	57187.478	13.92	15.13	16.44	18.17	2-m	VA
16.06.2015	57190.367	13.95	15.17	16.52	18.30	2-m	VA
16.07.2015	57220.364	13.91	15.26	16.44	18.13	Sch	FLI
17.07.2015	57221.417	13.94	15.32	16.55	18.15	Sch	FLI
19.07.2015	57223.400	14.11	15.34	16.66	18.41	2-m	VA
20.07.2015	57224.422	14.20	15.45	16.83	18.61	2-m	VA
11.08.2015	57246.355	14.08	15.50	16.79	18.48	1.3-m	ANDOR
12.08.2015	57247.394	14.00	15.40	16.66	18.34	1.3-m	ANDOR

Table 13: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
17.08.2015	57252.384	14.05	15.25	16.56	18.34	2-m	VA
24.08.2015	57259.337	13.95	15.32	16.53	18.16	Sch	FLI
25.08.2015	57260.342	14.06	15.43	16.66	18.35	Sch	FLI
03.09.2015	57269.328	13.87	15.20	16.33	17.95	Sch	FLI
04.09.2015	57270.330	13.98	15.17	16.49	18.24	2-m	VA
05.09.2015	57271.323	13.91	15.09	16.39	18.13	2-m	VA
06.09.2015	57272.288	14.04	15.23	16.54	18.26	2-m	VA

Table 14: Photometric CCD observations and data of LkH α 191 during the period 2013 April–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
10.04.2013	56392.510	11.56	12.14	12.79	13.87	Sch	FLI
11.04.2013	56394.482	11.56	12.16	12.83	13.91	Sch	FLI
02.05.2013	56415.418	11.56	12.17	12.82	13.96	Sch	FLI
15.05.2013	56428.404	11.58	12.15	12.87	13.92	60-cm	FLI
17.05.2013	56430.408	11.59	12.21	12.88	13.94	60-cm	FLI
19.05.2013	56432.405	11.56	12.17	12.87	13.90	60-cm	FLI
30.05.2013	56443.367	11.57	12.16	12.84	13.94	Sch	FLI
31.05.2013	56444.354	11.57	12.15	12.83	13.94	Sch	FLI
04.08.2013	56509.288	11.57	12.17	12.88	13.96	Sch	FLI
05.08.2013	56510.369	11.56	12.16	12.88	13.93	60-cm	FLI
05.08.2013	56510.386	11.56	12.16	12.83	13.95	Sch	FLI
06.08.2013	56511.411	11.58	12.17	12.85	13.96	Sch	FLI
06.08.2013	56511.413	11.56	12.19	12.86	13.95	60-cm	FLI
07.08.2013	56512.398	11.54	12.12	12.80	13.90	Sch	FLI
07.08.2013	56512.403	11.51	12.10	12.82	13.89	60-cm	FLI
08.08.2013	56513.382	11.53	12.17	12.84	14.01	60-cm	FLI
09.08.2013	56514.350	11.53	12.14	12.86	13.92	60-cm	FLI
12.08.2013	56517.284	11.53	12.11	12.85	13.94	60-cm	FLI
04.09.2013	56540.274	11.59	12.18	12.85	13.99	Sch	FLI
05.09.2013	56541.322	11.57	12.17	12.84	13.93	Sch	FLI
06.09.2013	56542.381	11.58	12.19	12.85	13.94	Sch	FLI
11.09.2013	56547.383	11.56	12.16	12.86	13.96	60-cm	FLI
14.09.2013	56550.363	11.59	12.18	12.89	14.00	60-cm	FLI
12.10.2013	56578.331	11.48	12.13	12.80	14.03	60-cm	FLI
28.12.2013	56655.200	11.55	12.14	12.82	13.90	Sch	FLI
29.12.2013	56656.180	11.57	12.16	12.84	13.94	Sch	FLI
30.12.2013	56657.193	11.56	12.13	12.82	13.92	Sch	FLI
23.01.2014	56681.189	11.57	12.16	12.83	13.96	Sch	FLI
22.03.2014	56738.590	11.56	12.18	12.85	13.96	Sch	FLI
21.05.2014	56799.425	11.58	12.16	12.84	13.96	Sch	FLI
28.06.2014	56837.417	11.58	12.18	12.85	13.95	Sch	FLI
29.06.2014	56838.398	11.57	12.19	12.86	13.94	Sch	FLI
21.07.2014	56860.391	11.50	12.16	12.79	13.93	60-cm	FLI
24.07.2014	56863.339	11.58	12.18	-	13.98	Sch	FLI
25.07.2014	56864.356	11.59	12.21	-	-	Sch	FLI
03.08.2014	56873.386	11.56	12.16	12.84	13.95	Sch	FLI
18.08.2014	56888.450	11.58	12.19	12.85	13.96	Sch	FLI
19.08.2014	56889.365	11.56	12.17	12.86	13.96	Sch	FLI
26.11.2014	56988.181	11.60	12.19	12.87	13.98	Sch	FLI
13.12.2014	57005.181	11.54	12.17	12.86	13.96	Sch	FLI
14.12.2014	57006.220	11.58	12.20	12.92	-	Sch	FLI

Table 14: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
21.02.2015	57074.641	11.61	12.22	12.93	14.05	Sch	FLI
23.04.2015	57136.487	11.59	12.24	12.92	14.04	Sch	FLI
25.04.2015	57138.491	11.61	12.24	12.92	14.04	Sch	FLI
18.05.2015	57161.423	11.61	12.22	12.93	14.04	Sch	FLI
19.05.2015	57162.460	11.60	12.23	12.92	14.06	Sch	FLI
21.05.2015	57164.459	11.62	12.25	12.93	14.06	Sch	FLI
12.06.2015	57186.444	11.60	12.23	12.97	14.04	Sch	FLI
16.07.2015	57220.364	11.60	12.23	12.87	14.03	Sch	FLI
17.07.2015	57221.417	11.60	12.22	12.90	14.05	Sch	FLI
24.08.2015	57259.337	11.60	12.23	12.94	14.04	Sch	FLI
25.08.2015	57260.342	11.59	12.23	12.93	14.04	Sch	FLI
03.09.2015	57269.328	11.58	12.21	12.93	14.01	Sch	FLI

Table 15: Photometric CCD observations and data of [KW97] 53-11 during the period 2013 April–2015 September

Date	J.D. (24...)	I	R	Tel	CCD
10.04.2013	56392.510	16.11	-	Sch	FLI
11.04.2013	56394.482	16.09	-	Sch	FLI
02.05.2013	56415.418	16.19	-	Sch	FLI
15.05.2013	56428.404	16.26	-	60-cm	FLI
17.05.2013	56430.408	16.31	-	60-cm	FLI
19.05.2013	56432.405	16.30	-	60-cm	FLI
30.05.2013	56443.367	16.10	-	Sch	FLI
31.05.2013	56444.354	16.22	-	Sch	FLI
01.08.2013	56506.281	16.09	18.61	2-m	VA
02.08.2013	56507.272	16.14	19.02	2-m	VA
03.08.2013	56508.318	16.11	18.64	2-m	VA
04.08.2013	56509.288	16.27	-	Sch	FLI
05.08.2013	56510.369	16.11	-	60-cm	FLI
05.08.2013	56510.386	16.23	-	Sch	FLI
06.08.2013	56511.411	16.21	-	Sch	FLI
06.08.2013	56511.413	16.26	-	60-cm	FLI
07.08.2013	56512.398	16.27	-	Sch	FLI
07.08.2013	56512.403	16.13	-	60-cm	FLI
08.08.2013	56513.382	16.23	-	60-cm	FLI
09.08.2013	56514.350	16.17	-	60-cm	FLI
12.08.2013	56517.284	16.31	-	60-cm	FLI
04.09.2013	56540.274	16.25	-	Sch	FLI
05.09.2013	56541.322	16.18	-	Sch	FLI
06.09.2013	56542.381	16.23	-	Sch	FLI
11.09.2013	56547.383	16.23	-	60-cm	FLI
14.09.2013	56550.363	16.22	-	60-cm	FLI
17.09.2013	56553.282	16.08	-	1.3-m	ANDOR
11.10.2013	56577.306	16.18	-	60-cm	FLI
12.10.2013	56578.331	16.22	-	60-cm	FLI
07.11.2013	56604.264	16.22	-	60-cm	FLI
09.12.2013	56636.185	15.84	18.91	2-m	VA
28.12.2013	56655.200	16.19	-	Sch	FLI
29.12.2013	56656.180	16.20	-	Sch	FLI
30.12.2013	56657.193	16.26	-	Sch	FLI
23.01.2014	56681.189	16.19	-	Sch	FLI
22.03.2014	56738.590	16.22	-	Sch	FLI

Table 15: Continued.

Date	J.D. (24...)	I	R	Tel	CCD
21.05.2014	56799.425	16.13	-	Sch	FLI
23.05.2014	56801.436	16.18	18.71	2-m	VA
23.06.2014	56832.391	16.15	18.72	2-m	VA
25.06.2014	56834.357	-	18.64	2-m	VA
26.06.2014	56835.462	16.14	18.68	2-m	VA
28.06.2014	56837.417	16.24	-	Sch	FLI
29.06.2014	56838.398	16.14	-	Sch	FLI
20.07.2014	56859.390	16.09	-	60-cm	FLI
21.07.2014	56860.391	16.13	-	60-cm	FLI
24.07.2014	56863.339	16.13	-	Sch	FLI
25.07.2014	56864.356	16.08	-	Sch	FLI
03.08.2014	56873.386	16.21	-	Sch	FLI
04.08.2014	56874.407	16.18	-	Sch	FLI
18.08.2014	56888.450	16.11	-	Sch	FLI
19.08.2014	56889.365	16.19	-	Sch	FLI
29.08.2014	56899.303	16.15	-	1.3-m	ANDOR
26.11.2014	56988.181	16.22	-	Sch	FLI
13.12.2014	57005.181	16.18	-	Sch	FLI
14.12.2014	57006.220	16.21	-	Sch	FLI
24.12.2014	57016.174	16.09	18.68	2-m	VA
25.12.2014	57017.175	16.11	18.72	2-m	VA
21.02.2015	57074.641	16.21	-	Sch	FLI
23.04.2015	57136.487	16.29	-	Sch	FLI
25.04.2015	57138.491	16.18	-	Sch	FLI
18.05.2015	57161.423	16.26	-	Sch	FLI
19.05.2015	57162.460	16.13	-	Sch	FLI
21.05.2015	57164.459	16.24	-	Sch	FLI
24.05.2015	57167.409	16.17	18.48	2-m	VA
12.06.2015	57186.444	16.25	-	Sch	FLI
13.06.2015	57187.478	16.04	18.47	2-m	VA
16.06.2015	57190.367	16.06	18.51	2-m	VA
16.07.2015	57220.364	16.15	-	Sch	FLI
17.07.2015	57221.417	16.25	-	Sch	FLI
19.07.2015	57223.400	16.11	18.50	2-m	VA
20.07.2015	57224.422	16.12	18.50	2-m	VA
11.08.2015	57246.355	16.11	19.91	1.3-m	ANDOR
12.08.2015	57247.394	16.11	19.88	1.3-m	ANDOR
24.08.2015	57259.337	16.19	-	Sch	FLI
25.08.2015	57260.342	16.21	-	Sch	FLI
03.09.2015	57269.328	16.22	-	Sch	FLI
04.09.2015	57270.330	16.12	18.71	2-m	VA
05.09.2015	57271.323	16.17	18.58	2-m	VA
06.09.2015	57272.288	16.16	18.52	2-m	VA

Table 16: Photometric CCD observations and data of [KW97] 53-23 during the period 2013 April–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
10.04.2013	56392.510	14.98	-	-	-	Sch	FLI
11.04.2013	56394.482	14.92	17.30	-	-	Sch	FLI
02.05.2013	56415.418	14.99	17.29	-	-	Sch	FLI
15.05.2013	56428.404	15.05	17.25	-	-	60-cm	FLI
17.05.2013	56430.408	14.95	17.30	-	-	60-cm	FLI

Table 16: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
19.05.2013	56432.405	15.10	16.95	-	-	60-cm	FLI
30.05.2013	56443.367	14.98	17.43	-	-	Sch	FLI
31.05.2013	56444.354	14.88	17.36	-	-	Sch	FLI
04.07.2013	56478.363	15.24	16.82	18.68	-	2-m	VA
01.08.2013	56506.281	15.22	16.88	18.76	-	2-m	VA
02.08.2013	56507.272	15.09	16.69	18.49	-	2-m	VA
03.08.2013	56508.318	15.12	16.76	18.62	-	2-m	VA
04.08.2013	56509.288	14.94	17.19	19.06	-	Sch	FLI
05.08.2013	56510.369	14.99	17.02	-	-	60-cm	FLI
05.08.2013	56510.386	14.87	16.99	18.93	-	Sch	FLI
06.08.2013	56511.411	14.90	17.07	18.85	-	Sch	FLI
06.08.2013	56511.413	15.04	17.02	18.54	-	60-cm	FLI
07.08.2013	56512.398	14.86	17.04	18.84	-	Sch	FLI
07.08.2013	56512.403	14.98	17.04	18.73	-	60-cm	FLI
08.08.2013	56513.382	14.97	17.02	18.76	-	60-cm	FLI
09.08.2013	56514.350	14.99	17.00	18.71	-	60-cm	FLI
12.08.2013	56517.284	14.94	16.99	-	-	60-cm	FLI
04.09.2013	56540.274	14.97	17.13	19.01	-	Sch	FLI
05.09.2013	56541.322	14.90	17.08	18.66	-	Sch	FLI
06.09.2013	56542.381	14.95	17.10	18.62	-	Sch	FLI
07.09.2013	56543.396	15.09	16.68	18.51	-	2-m	VA
08.09.2013	56544.264	15.07	16.67	18.42	-	2-m	VA
11.09.2013	56547.383	14.96	17.01	-	-	60-cm	FLI
14.09.2013	56550.363	14.95	17.21	18.82	-	60-cm	FLI
17.09.2013	56553.282	14.92	16.91	18.55	-	1.3-m	ANDOR
11.10.2013	56577.306	-	-	-	-	60-cm	FLI
12.10.2013	56578.331	-	-	-	-	60-cm	FLI
07.11.2013	56604.264	15.16	17.48	-	-	60-cm	FLI
09.12.2013	56636.185	15.34	17.01	18.94	-	2-m	VA
28.12.2013	56655.200	15.19	-	-	-	Sch	FLI
29.12.2013	56656.180	15.27	17.58	19.30	-	Sch	FLI
30.12.2013	56657.193	15.06	17.43	19.14	-	Sch	FLI
23.01.2014	56681.189	15.09	17.39	19.19	-	Sch	FLI
05.02.2014	56694.194	-	17.02	19.04	-	2-m	VA
22.03.2014	56738.590	15.07	17.51	19.27	-	Sch	FLI
21.05.2014	56799.425	15.24	17.61	-	-	Sch	FLI
23.05.2014	56801.436	15.31	16.94	18.79	-	2-m	VA
23.06.2014	56832.391	15.15	16.74	18.57	-	2-m	VA
25.06.2014	56834.357	15.20	16.87	18.83	-	2-m	VA
26.06.2014	56835.462	15.29	16.91	18.84	-	2-m	VA
28.06.2014	56837.417	15.10	17.43	19.05	-	Sch	FLI
29.06.2014	56838.398	15.21	17.60	19.49	-	Sch	FLI
20.07.2014	56859.390	14.95	16.95	-	-	60-cm	FLI
21.07.2014	56860.391	14.95	16.95	-	-	60-cm	FLI
24.07.2014	56863.339	15.01	17.17	-	-	Sch	FLI
25.07.2014	56864.356	15.04	17.47	-	-	Sch	FLI
03.08.2014	56873.386	15.00	17.29	19.10	-	Sch	FLI
04.08.2014	56874.407	15.17	17.73	-	-	Sch	FLI
18.08.2014	56888.450	15.15	17.57	-	-	Sch	FLI
19.08.2014	56889.365	15.04	17.36	19.21	-	Sch	FLI
29.08.2014	56899.303	15.21	17.20	18.98	-	1.3-m	ANDOR
26.11.2014	56988.181	15.06	17.46	18.90	-	Sch	FLI
13.12.2014	57005.181	15.12	17.58	18.89	-	Sch	FLI
14.12.2014	57006.220	15.15	17.56	19.40	-	Sch	FLI
24.12.2014	57016.174	15.16	16.84	18.75	-	2-m	VA
25.12.2014	57017.175	15.30	16.98	18.86	-	2-m	VA
21.02.2015	57074.641	15.02	17.08	18.72	-	Sch	FLI

Table 16: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
23.04.2015	57136.487	15.06	17.07	18.78	-	Sch	FLI
25.04.2015	57138.491	14.95	16.96	18.61	-	Sch	FLI
18.05.2015	57161.423	15.14	17.06	-	-	Sch	FLI
19.05.2015	57162.460	14.98	16.91	-	-	Sch	FLI
21.05.2015	57164.459	15.07	17.04	18.61	-	Sch	FLI
24.05.2015	57167.409	15.00	16.58	18.41	20.39	2-m	VA
12.06.2015	57186.444	15.05	17.01	18.58	-	Sch	FLI
13.06.2015	57187.478	15.20	16.81	18.77	20.24	2-m	VA
16.06.2015	57190.367	15.09	16.68	18.56	-	2-m	VA
16.07.2015	57220.364	14.98	16.97	18.51	-	Sch	FLI
17.07.2015	57221.417	14.98	17.00	18.46	-	Sch	FLI
19.07.2015	57223.400	14.97	16.60	18.39	20.49	2-m	VA
20.07.2015	57224.422	14.96	16.49	18.32	20.41	2-m	VA
11.08.2015	57246.355	15.06	17.03	18.70	-	1.3-m	ANDOR
12.08.2015	57247.394	14.98	16.96	18.57	-	1.3-m	ANDOR
17.08.2015	57252.384	15.18	16.78	18.55	-	2-m	VA
24.08.2015	57259.337	15.07	17.00	-	-	Sch	FLI
25.08.2015	57260.342	15.09	17.14	18.58	-	Sch	FLI
03.09.2015	57269.328	15.20	17.22	19.01	-	Sch	FLI
04.09.2015	57270.330	15.19	16.84	18.84	-	2-m	VA
05.09.2015	57271.323	15.10	16.66	18.37	20.24	2-m	VA
06.09.2015	57272.288	15.12	16.66	18.44	-	2-m	VA

Table 17: Photometric CCD observations and data of [KW97] 53-36 during the period 2013 April–2015 September

Date	J.D. (24...)	I	R	V	B	Tel	CCD
10.04.2013	56392.510	11.61	11.86	12.28	12.99	Sch	FLI
11.04.2013	56394.482	11.61	11.87	12.28	13.04	Sch	FLI
02.05.2013	56415.418	11.63	11.87	12.26	13.05	Sch	FLI
15.05.2013	56428.404	11.63	11.90	12.29	13.04	60-cm	FLI
17.05.2013	56430.408	11.65	11.91	12.33	13.05	60-cm	FLI
30.05.2013	56443.367	11.62	11.84	12.29	13.06	Sch	FLI
31.05.2013	56444.354	11.62	11.85	12.26	13.05	Sch	FLI
04.08.2013	56509.288	11.61	11.86	12.29	13.05	Sch	FLI
05.08.2013	56510.369	11.63	11.90	12.32	13.03	60-cm	FLI
05.08.2013	56510.386	11.61	11.86	12.26	13.06	Sch	FLI
06.08.2013	56511.411	11.62	11.86	12.26	13.05	Sch	FLI
06.08.2013	56511.413	11.62	11.89	12.29	13.02	60-cm	FLI
07.08.2013	56512.398	11.62	11.87	12.28	13.06	Sch	FLI
07.08.2013	56512.403	11.61	11.86	12.30	13.00	60-cm	FLI
08.08.2013	56513.382	11.59	11.88	12.28	13.07	60-cm	FLI
09.08.2013	56514.350	11.60	11.89	12.31	13.01	60-cm	FLI
12.08.2013	56517.284	11.58	11.84	12.27	12.99	60-cm	FLI
04.09.2013	56540.274	11.62	11.86	12.26	13.05	Sch	FLI
05.09.2013	56541.322	11.61	11.86	12.27	13.02	Sch	FLI
06.09.2013	56542.381	11.62	11.88	12.28	13.02	Sch	FLI
11.09.2013	56547.383	11.62	11.88	12.30	13.03	60-cm	FLI
14.09.2013	56550.363	11.66	11.91	12.33	13.08	60-cm	FLI
28.12.2013	56655.200	11.61	11.86	12.26	13.02	Sch	FLI
29.12.2013	56656.180	11.63	11.87	12.27	13.04	Sch	FLI
30.12.2013	56657.193	11.62	11.86	12.27	13.03	Sch	FLI
23.01.2014	56681.189	11.62	11.86	12.27	13.06	Sch	FLI

Table 17: Continued.

Date	J.D. (24...)	I	R	V	B	Tel	CCD
22.03.2014	56738.590	11.63	11.88	12.28	13.07	Sch	FLI
21.05.2014	56799.425	11.63	11.86	12.27	13.07	Sch	FLI
28.06.2014	56837.417	11.62	11.86	12.27	13.03	Sch	FLI
29.06.2014	56838.398	11.62	11.88	12.27	13.01	Sch	FLI
24.07.2014	56863.339	11.63	11.90	-	13.06	Sch	FLI
25.07.2014	56864.356	11.64	11.90	-	-	Sch	FLI
03.08.2014	56873.386	11.62	11.86	12.27	13.04	Sch	FLI
18.08.2014	56888.450	11.63	11.87	12.27	13.03	Sch	FLI
19.08.2014	56889.365	11.62	11.86	12.28	13.05	Sch	FLI
26.11.2014	56988.181	11.65	11.88	12.26	13.06	Sch	FLI
13.12.2014	57005.181	11.53	11.85	12.17	12.99	Sch	FLI
14.12.2014	57006.220	11.60	11.90	12.19	-	Sch	FLI
21.02.2015	57074.641	11.63	11.91	12.25	13.13	Sch	FLI
23.04.2015	57136.487	11.61	11.90	12.26	13.08	Sch	FLI
25.04.2015	57138.491	11.62	11.90	12.30	13.10	Sch	FLI
19.05.2015	57162.460	11.62	11.89	12.26	13.10	Sch	FLI
12.06.2015	57186.444	11.62	11.90	12.24	13.09	Sch	FLI
16.07.2015	57220.364	11.62	11.92	12.28	13.12	Sch	FLI
17.07.2015	57221.417	11.63	11.93	12.33	13.17	Sch	FLI
24.08.2015	57259.337	11.65	11.96	12.28	13.16	Sch	FLI
25.08.2015	57260.342	11.63	11.92	-	13.12	Sch	FLI
03.09.2015	57269.328	11.64	11.92	-	13.10	Sch	FLI